McGirr, R., Seton, M., and Williams, S., 2020, Kinematic and geodynamic evolution of the Isthmus of Panama region: Implications for Central American Seaway closure: GSA Bulletin, <https://doi.org/10.1130/B35595.1>.

Supplemental Material

Figure S1. Müller et al. (2016) kinematic model and corresponding paleo-age-grid from 35 to present-day. Reconstructed continental crust (dark-gray), coastlines (light-gray), subduction zones (black lines with filled teeth), ridges (black lines), extinct ridges (thin black lines), topology network polygons (white and dark gray lines) and isochrons (thin gray lines). Caribbean (Car); East Pacific Rise (EPR); Pacific (Pac); Cocos-Nazca Spreading (CNS); South America (SAM); Bauer Microplate (BM); North America (NAM); Mathematician (M).

Figure S2. Shephard et al. (2013) kinematic model and corresponding paleo-age-grid from 35 to present-day. Abbreviations and lines the same as Fig. S1.

Figure S3. Tomographic profiles from Obayashi et al. (2013) global P-wave tomography model sampled from profiles shown in Figure 8. Color indicates anomalies in seismic wave speed as a $\pm$ 0.8% deviation from average mantle velocities, indicative of subducted slab material (blue), or regions of upwelling (red).

Figure S4. Tomographic profiles from Montelli et al. (2006) global P-wave tomography model sampled from profiles shown in Figure 8. See Fig. S3 for interpretation.

Figure S5. Tomographic profiles from Grand (2002) global S-wave tomography model sampled from profiles shown in Figure 8. See Fig. S3 for interpretation.

Table S1. Alternative global P- and S-wave seismic tomography models analyzed beneath Central and northern South America.

The preferred plate motion model and slab-tracker code which was used to generate reconstructions of slab geometries using GPlates-format topological plate reconstructions is available at the following GitHub repository: <https://github.com/siwill22/pgpslabs>.

Magnetic anomaly data

The plate reconstruction required the compilation of previously digitised east Pacific Ocean magnetic anomaly identifications (Seton et al., 2014). Magnetic anomaly “picks” define the location where the ship track of a magnetic survey intersects an anomaly. These picks are assigned a chron number and “chron end”. Each pick in the region had previously been assigned GPlates Mark-up Language (GPML) attributes, including appearance and disappearance ages, according to the timescale of Gee and Kent (2007), and were reassigned Plate Identification Numbers (PIDs) based on their present-day location within the plate polygons associated with Matthews et al. (2016). New PIDs were assigned where magnetic picks were associated with fossil spreading centers (MacLeod et al., 2017), not yet implemented into the most recent plate reconstruction. Magnetic anomaly data (Fig. 3) were the primary constraint for our improved plate model.

Bathymetric features

In addition to our preferred magnetic anomaly picks, we visually reconstructed the Panama Basin by identifying the origin and age of notable bathymetric features (Fig. 3), derived from satellite altimetry (Sandwell and Smith, 2009). The age and origin of aseismic ridges resulting from mantle plume upwellings in the region were used to inform direction and rate of seafloor spreading (Hey, 1977; Lonsdale and Klitgord, 1978) and the location and timing of ridge jumps (Morell, 2015). Other bathymetric features in the region such as fracture zones (Matthews et al., 2016) and extinct ridges (MacLeod et al., 2017) gave an indication to the past location of plate boundaries, and defined the direction of seafloor-spreading.

Rotation model

Our kinematic reconstruction required the transformation of the geological observations explained previously into a quantitative framework, which we constructed using GPlates (Müller et al., 2018). We defined the relative motions of plates via a set of finite rotations, which are the total rotation pole of a plate with respect to its present-day position as defined by Cox and Hart (1986), and their corresponding Euler poles (Müller et al., 2018). The calculation of Euler pole locations and angles of rotation required the digitisation of magnetic lineations guided by our set of preferred magnetic anomaly picks and fracture zone crossings. Magnetic lineations were visually paired with conjugate segments on the conjugate plate of a spreading system. Spreading direction was guided by small circles that describe the motion between Cocos and Nazca plates, and Galapagos plume products that trace Cocos and Nazca absolute plate motion. Additionally, we implemented a modified version of the Oligocene to present model of Montes et al. (2012) for Central American arc formation, based on paleomagnetic analysis (Di Marco et al., 1995).

Seafloor spreading agegrids

The updated global rotation model and seafloor-spreading isochrons were used to calculate continuous grids of global age-area distribution of oceanic crust from 230 Ma to present, at million year intervals. The grids were computed by deriving a set of densely-interpolated seafloor-spreading isochrons and gridding in GMT (Müller et al., 2008). The age-gridding process required additional inputs that included the location of present-day mid ocean ridges, extinct ridges, and isochrons defining the age of continent-ocean boundaries (COBs). The paleo-agegrids were masked with COB polygons that define areas of continental crust or crust of unknown origin.

Topologically closed plate networks

We utilized the approach developed by Gurnis et al. (2012) to create a self-consistent network of dynamically closing plate polygons on Earth's surface within GPlates. Each plate polygon is composed of a finite list of plate margins, all with which are defined by an individual Euler pole. The continuously closing plate (CCP) method ensures that as each feature forming the boundary of a plate polygon moves independently in the reconstruction, the plate polygon remains topologically closed as a function of time (Gurnis et al., 2012). We have improved the dynamic plate polygon and plate boundary data set for the Oligocene to recent based on magnetic anomaly lineations in the eastern Pacific Ocean, previous work on the timing of Panama-South America collision (Montes et al., 2012), and the geometry of subducted segments of the Cocos-Nazca system (Johnston and Thorkelson, 1997), based on geological observations.

Shallow-dipping slabs

We have included the evolution of the Colombian flat slab within our tectonic reconstruction following the approach of Flament et al. (2015). The evolution of the extent of the flat slab segment is based on geological constraints, subduction zone geometries, and the velocity of the Nazca plate. A scenario was imposed in which the flat slab progressively shallows from 9 Ma, in accordance with the model of Colombian flat slab evolution of Wagner et al. (2017) (Fig. 2C), by redefining the depth of the flat slab every million years until a persistent flat slab is acquired at 6 Ma. The maximum depth of the subduction zone was redefined every million years to ensure that the outboard trench of the flat slab segment is not deeper than the flat slab itself following Equation (1);

$$Z_{SZ} = Z_{FS} + 30A_{FS}, \quad (1)$$

where Z_{SZ} is subduction zone depth in km, Z_{FS} is flat slab depth in km, and A_{FS} is the age of the flat slab in Myr. This assumes that the leading edge of the subducting slab sinks in the upper mantle at a rate of 30 mm yr^{-1} (van der Meer et al., 2010).

We used the method of Gurnis et al. (2012) to define the regional extent of the Colombian flat slab in GPlates by dynamically closing the flat slab polygon, which was defined by the leading edge of the flat slab and the northernmost segment of the South American Trench (SAT). The southern boundary of the flat slab segment was defined by the present-day location of the Caldas tear (a Wadati-Benioff zone offset) which coincides with the location of the

subducting, inactive Sandra Rift (Lonsdale, 2005; Chiarabba et al., 2016; Wagner et al., 2017). The flat slab segment extends farther north than our model suggests, being fed by both Nazca and Caribbean subduction, and the contemporaneous underthrusting of the Panama arc (Idárraga-García et al., 2016). For simplicity and modeling limitations, we have only implemented the Nazca component.

Modeling subducted slab volumes

The methodology of Thorkelson (1996) defines slab window geometry according to the ridge-transform-trench configuration, slab dip, and plate motion vectors. We developed a program using PyGPlates that plots the surface projection of subduction volumes (and subsequently, slab gaps), by extracting the location of trench segments and plate motion vectors from alternative kinematic plate models (Fig. S1–2) (Shephard et al., 2013; Müller et al., 2016), including our revised global plate reconstruction, and applying an assumed slab dip (45°), based on published three-dimensional models of subduction zone geometries in the area of interest (Hayes et al., 2012).

The method involves extracting arcuate subduction trenches from the topological plate boundary model with common overriding and downgoing plates, then dividing these curved segments into shorter segments which are assumed to be straight and with equally spaced midpoints (~50 km apart). The stage pole that describes the motion between the overriding and downgoing plates were then used to calculate the velocity vector of each of the points, which were divided into trench orthogonal and perpendicular components. A new orthogonal velocity vector was calculated by correcting the orthogonal component for the assumed slab dip, which was then combined with the perpendicular vector to get the velocity and stage pole that describes the motion of each dipping slab point, following Equation (2);

$$\vec{V} = \overline{TP} + \cos(\theta)\overline{TN}, \quad (2)$$

where $\vec{V}$ is the new orthogonal velocity vector, $\overline{TP}$ is the trench parallel vector, θ is the dip angle of the subducting slab, and $\overline{TN}$ is the trench normal vector. The new vector and stage pole were then used to reconstruct the points along the trench at 1 Myr intervals, and get the increase in depth of each point, following Equation (3);

$$\Delta z = \sin(\theta)|\vec{V}|, \quad (3)$$

where Δz is the change in depth and $\vec{V}$ is the velocity magnitude. This simple method ignores the effects of thermal erosion, spherical shell strain and upward curling of the window margins on window geometry (Thorkelson, 1996).

Finally, we extracted the age of oceanic crust for each point at the time of trench intersection from a set of paleo-age-grids, corresponding to the associated kinematic model. This was repeated for a series of timesteps, between 30 and 15 Ma, and 15 and 0 Ma. We visualized the results by plotting a snapshot of subducted slab points in the upper mantle (depths <700 km) incrementally reconstructed from topological plate configurations within each period (Fig. 7), constrained by the input kinematic model for the Panama Isthmus region. Slab windows are visualized as the gaps between the reconstructed slab segments.

Age-coded seismic tomography match to alternative kinematic models

We assumed that the age of subducted slab material can be inferred based on the relationship between sinking rate and mantle viscosity, such that slab material will sink slower through a more viscous medium. We apply an age-depth conversion to depth-dependent rasters of various seismic tomography models given a sinking rate of 4.8 cm yr⁻¹ for the less viscous upper mantle (Lithgow-Bertelloni and Richards, 1998), and a sinking rate of 1.2 cm yr⁻¹ for the viscous lower mantle (van der Meer et al., 2010).

Horizontal depth slices for each analyzed seismic tomography model were age-coded using the upper and lower mantle sinking rates defined above, following Equation (4);

$$A = \begin{cases} \frac{z}{v_{um}} & \text{where } z \leq 660 \text{ km} \\ \frac{660}{v_{um}} + \frac{660-z}{v_{lm}} & \text{where } 660 \text{ km} < z \leq 2890 \text{ km} \end{cases} \quad (4)$$

where A is the age of the subducted slab material in Myrs, z is the depth of the horizontal tomography slice in km, v_{um} and v_{lm} are the sinking velocities in mm yr⁻¹, applied to the upper and lower mantle respectively.

It is necessary to note that the age-depth conversion is relevant to slab volumes only, and is not applicable to the adjacent mantle. The age assignments are an estimate, and do not account for variations in sinking rate which may vary regionally depending on plate convergence rates, amongst other subduction parameters.

REFERENCES CITED

- Chiarabba, C., De Gori, P., Faccenna, C., Speranza, F., Seccia, D., Dionicio, V., and Prieto, G.A., 2016, Subduction system and flat slab beneath the Eastern Cordillera of Colombia: *Geochemistry Geophysics Geosystems*, v. 17, no. 1, p. 16–27, <https://doi.org/10.1002/2015GC006048>.
- Cox, A., and Hart, B., 1986, *Plate tectonics: How it works*: Palo Alto, Blackwell Scientific Publications, 392 p.
- Di Marco, G., Baumgartner, P.O., and Channell, J.E.T., 1995, Late Cretaceous-early Tertiary paleomagnetic data and a revised tectonostratigraphic subdivision of Costa Rica and western Panama, *in* Mann, P., ed., *Geologic and tectonic development of the Caribbean plate boundary in southern Central America*: Boulder, Colorado, Geological Society of America Special Paper 295, p. 1–28, <https://doi.org/10.1130/SPE295-p1>.
- Flament, N., Gurnis, M., Müller, R.D., Bower, D.J., and Husson, L., 2015, Influence of subduction history on South American topography: *Earth and Planetary Science Letters*, v. 430, p. 9–18, <https://doi.org/10.1016/j.epsl.2015.08.006>.
- Gee, J.S., and Kent, D.V., 2007, Source of oceanic magnetic anomalies and the geomagnetic polarity time scale: *Treatise on Geophysics*, v. 5, p. 455–507, <https://doi.org/10.1016/B978-04452748-6/00097-3>.
- Grand, S.P., 2002, Mantle shear-wave tomography and the fate of subducted slabs: *Philosophical Transactions of the Royal Society of London, Series A: Mathematical, Physical and Engineering Sciences*, v. 360, no. 1800, p. 2475–2491, <https://doi.org/10.1098/rsta.2002.1077>.

- Gurnis, M., Turner, M., Zahirovic, S., DiCaprio, L., Spasojevic, S., Müller, R.D., Boyden, J., Seton, M., Manea, V.C., and Bower, D.J., 2012, Plate tectonic reconstructions with continuously closing plates: *Computers & Geosciences*, v. 38, no. 1, p. 35–42, <https://doi.org/10.1016/j.cageo.2011.04.014>.
- Hayes, G.P., Wald, D.J., and Johnson, R.L., 2012, Slab1.0: A three-dimensional model of global subduction zone geometries: *Journal of Geophysical Research. Solid Earth*, v. 117, no. B1, <https://doi.org/10.1029/2011JB008524>.
- Hey, R., 1977, Tectonic evolution of the Cocos-Nazca spreading center: *Geological Society of America Bulletin*, v. 88, no. 12, p. i–vi, [https://doi.org/10.1130/0016-7606\(1977\)88<i:TEOTCS>2.0.CO;2](https://doi.org/10.1130/0016-7606(1977)88<i:TEOTCS>2.0.CO;2).
- Idárraga-García, J., Kendall, J.M., and Vargas, C.A., 2016, Shear wave anisotropy in northwestern South America and its link to the Caribbean and Nazca subduction geodynamics: *Geochemistry Geophysics Geosystems*, v. 17, no. 9, p. 3655–3673, <https://doi.org/10.1002/2016GC006323>.
- Johnston, S.T., and Thorkelson, D.J., 1997, Cocos-Nazca slab window beneath central America: *Earth and Planetary Science Letters*, v. 146, no. 3–4, p. 465–474, [https://doi.org/10.1016/S0012-821X\(96\)00242-7](https://doi.org/10.1016/S0012-821X(96)00242-7).
- Li, C., van der Hilst, R.D., Engdahl, E.R., and Burdick, S., 2008, A new global model for P wave speed variations in Earth's mantle: *Geochemistry Geophysics Geosystems*, v. 9, no. 5, <https://doi.org/10.1029/2007GC001806>.
- Lithgow-Bertelloni, C., and Richards, M.A., 1998, The dynamics of Cenozoic and Mesozoic plate motions: *Reviews of Geophysics*, v. 36, no. 1, p. 27–78, <https://doi.org/10.1029/97RG02282>.
- Lonsdale, P., and Klitgord, K.D., 1978, Structure and tectonic history of the eastern Panama Basin: *Geological Society of America Bulletin*, v. 89, no. 7, p. 981–999, [https://doi.org/10.1130/0016-7606\(1978\)89<981:SATHOT>2.0.CO;2](https://doi.org/10.1130/0016-7606(1978)89<981:SATHOT>2.0.CO;2).
- Lonsdale, P., 2005, Creation of the Cocos and Nazca plates by fission of the Farallon plate: *Tectonophysics*, v. 404, no. 3–4, p. 237–264, <https://doi.org/10.1016/j.tecto.2005.05.011>.
- MacLeod, S.J., Williams, S.E., Matthews, K.J., Müller, R.D., and Qin, X., 2017, A global review and digital database of large-scale extinct spreading centers: *Geosphere*, v. 13, no. 3, p. 911–949, <https://doi.org/10.1130/GES01379.1>.
- Matthews, K.J., Maloney, K.T., Zahirovic, S., Williams, S.E., Seton, M., and Mueller, R.D., 2016, Global plate boundary evolution and kinematics since the late Paleozoic: *Global and Planetary Change*, v. 146, p. 226–250, <https://doi.org/10.1016/j.gloplacha.2016.10.002>.
- Montelli, R., Nolet, G., Dahlen, F.A., and Masters, G., 2006, A catalogue of deep mantle plumes: New results from finite-frequency tomography: *Geochemistry Geophysics Geosystems*, v. 7, no. 11, <https://doi.org/10.1029/2006GC001248>.
- Montes, C., Bayona, G., Cardona, A., Buchs, D.M., Silva, C.A., Morón, S., Hoyos, N., Ramírez, D.A., Jaramillo, C.A., and Valencia, V., 2012, Arc-continent collision and orocline formation: Closing of the Central American seaway: *Journal of Geophysical Research. Solid Earth*, v. 117, no. B4, <https://doi.org/10.1029/2011JB008959>.
- Morell, K.D., 2015, Late Miocene to recent plate tectonic history of the southern Central America convergent margin: *Geochemistry Geophysics Geosystems*, v. 16, no. 10, p. 3362–3382, <https://doi.org/10.1002/2015GC005971>.

- Müller, R.D., Sdrolias, M., Gaina, C., and Roest, W.R., 2008, Age, spreading rates, and spreading asymmetry of the world's ocean crust: *Geochemistry Geophysics Geosystems*, v. 9, no. 4, <https://doi.org/10.1029/2007GC001743>.
- Müller, R.D., Seton, M., Zahirovic, S., Williams, S.E., Matthews, K.J., Wright, N.M., Shephard, G.E., Maloney, K.T., Barnett-Moore, N., Hosseinpour, M., Bower, D.J., and Cannon, J., 2016, Ocean basin evolution and global-scale plate reorganization events since Pangea breakup: *Annual Review of Earth and Planetary Sciences*, v. 44, p. 107–138, <https://doi.org/10.1146/annurev-earth-060115-012211>.
- Müller, R.D., Cannon, J., Qin, X., Watson, R.J., Gurnis, M., Williams, S., Pfaffelmoser, T., Seton, M., Russell, S.H., and Zahirovic, S., 2018, GPlates: Building a virtual Earth through deep time: *Geochemistry Geophysics Geosystems*, v. 19, no. 7, p. 2243–2261, <https://doi.org/10.1029/2018GC007584>.
- Obayashi, M., Yoshimitsu, J., Nolet, G., Fukao, Y., Shiobara, H., Sugioka, H., Miyamachi, H., and Gao, Y., 2013, Finite frequency whole mantle P wave tomography: Improvement of subducted slab images: *Geophysical Research Letters*, v. 40, no. 21, p. 5652–5657, <https://doi.org/10.1002/2013GL057401>.
- Sandwell, D.T., and Smith, W.H., 2009, Global marine gravity from retracked Geosat and ERS-1 altimetry: Ridge segmentation versus spreading rate: *Journal of Geophysical Research. Solid Earth*, v. 114, no. B1, <https://doi.org/10.1029/2008JB006008>.
- Seton, M., Whittaker, J.M., Wessel, P., Müller, R.D., DeMets, C., Merkouriev, S., Cande, S., Gaina, C., Eagles, G., Granot, R., and Stock, J., 2014, Community infrastructure and repository for marine magnetic identifications: *Geochemistry Geophysics Geosystems*, v. 15, no. 4, p. 1629–1641, <https://doi.org/10.1002/2013GC005176>.
- Shephard, G.E., Müller, R.D., and Seton, M., 2013, The tectonic evolution of the Arctic since Pangea breakup: Integrating constraints from surface geology and geophysics with mantle structure: *Earth-Science Reviews*, v. 124, p. 148–183, <https://doi.org/10.1016/j.earscirev.2013.05.012>.
- Thorkelson, D.J., 1996, Subduction of diverging plates and the principles of slab window formation: *Tectonophysics*, v. 255, no. 1–2, p. 47–63, [https://doi.org/10.1016/0040-1951\(95\)00106-9](https://doi.org/10.1016/0040-1951(95)00106-9).
- van der Meer, D.G., Spakman, W., van Hinsbergen, D.J., Amaru, M.L., and Torsvik, T.H., 2010, Towards absolute plate motions constrained by lower-mantle slab remnants: *Nature Geoscience*, v. 3, no. 1, p. 36–40, <https://doi.org/10.1038/ngeo708>.
- Wagner, L.S., Jaramillo, J.S., Ramírez-Hoyos, L.F., Monsalve, G., Cardona, A., and Becker, T.W., 2017, Transient slab flattening beneath Colombia: *Geophysical Research Letters*, v. 44, no. 13, p. 6616–6623, <https://doi.org/10.1002/2017GL073981>.

Figure S1. Müller et al. (2016) kinematic model and corresponding paleo-age-grid from 35 to present-day. Reconstructed continental crust (dark-gray), coastlines (light-gray), subduction zones (black lines with filled teeth), ridges (black lines), extinct ridges (thin black lines), topology network polygons (white and dark gray lines) and isochrons (thin gray lines). Caribbean (Car); East Pacific Rise (EPR); Pacific (Pac); Cocos-Nazca Spreading (CNS); South America (SAM); Bauer Microplate (BM); North America (NAM); Mathematician (M).

Figure S2. Shephard et al. (2013) kinematic model and corresponding paleo-age-grid from 35 to present-day. Abbreviations and lines the same as Fig. S1.

Table S1. Alternative global P- and S-wave seismic tomography models analyzed beneath Central and northern South America.

Model name	Model type	Reference
MIT-P08	Finite frequency mantle P-wave velocity	Li et al. (2008)
GAP P4	Finite frequency mantle P-wave velocity	Obayashi et al. (2013)
Montelli P	Finite frequency mantle P-wave velocity	Montelli et al. (2006)
Grand S	Mantle Shear wave velocity	Grand (2002)

Figure S3. Tomographic profiles from Obayashi et al. (2013) global P-wave tomography model sampled from profiles shown in Figure 8. Color indicates anomalies in seismic wave speed as a $\pm 0.8\%$ deviation from average mantle velocities, indicative of subducted slab material (blue), or regions of upwelling (red).

Figure S4. Tomographic profiles from Montelli et al. (2006) global P-wave tomography model sampled from profiles shown in Figure 8. See Fig. S3 for interpretation.

Figure S5. Tomographic profiles from Grand (2002) global S-wave tomography model sampled from profiles shown in Figure 8. See Fig. S3 for interpretation.