

ANALYTICAL METHODS

U-Pb dating

Samples and standards were acid-washed before being analysed to remove possible surface Pb contamination. The grains were analysed using a commercial LUV213 laser ablation system ($\lambda = 213\text{nm}$) (New Wave/MerchanTek), attached to a Hewlett Packard 4500s ICPMS (Jackson et al., 2004). All ablations were carried out in He. Ablation pits were about 50 μm in diameter. The time-resolved signals were processed using the GLITTER interactive software to select the portions of the grains that had suffered least lead loss, or gain of common Pb, and were thus closest to being concordant. U-Pb isotopic data results are given in Table 2.

The standard used in this work is the GEMOC-GJ-1 gem zircon (Elhlou et al., 2006), with a TIMS age of 608.5 Ma. This standard is run 2-4 times before and after each ten unknowns. Cross-analysis of other international standards also gives good results. Two analyses of Mud Tank zircon (734 ± 32 Ma; Black and Gulson, 1978) and two of zircon 91500 (1064 Ma; Wiedenbeck et al., 1995) were run during this work, and their mean values are within 1 s.d. of the recommended values.

U-Pb data for some well-characterised zircons

Zircon	TIMS 207/206	206/238	207/235	207/206
91500	1065.4			
This study n=9		1067 ± 3	1065 ± 4	1061 ± 8
Long-term work* n=83		1061 ± 36	1063 ± 29	1068 ± 26
Mud Tank	734 ± 32			
This study n=9		726 ± 5	729 ± 7	737 ± 16
Long-term work* n=73		733 ± 14	735 ± 8	739 ± 12

* Jackson et al., 2004

Hf-isotope analyses

Hf isotope analyses were carried out *in situ* with a UP 213 laser-ablation microprobe, attached to a Nu Plasma multi-collector ICPMS at GEMOC, Macquarie University. Most analyses were carried out with a beam diameter of ca 50 μm , a 4 Hz repetition rate, and energies of 0.1 mJ/pulse and 0.6 J/cm². Typical ablation times were 80-120 s, resulting in pits 40-60 μm deep. The methodology and analyses of standard solutions and standard zircons are described by Griffin et al. (2000).

The measured $^{176}\text{Lu}/^{177}\text{Hf}$ ratios are used to calculate initial $^{176}\text{Hf}/^{177}\text{Hf}$ ratios. The typical 2SE uncertainty on a single analysis of $^{176}\text{Lu}/^{177}\text{Hf}$ is $\pm 1\text{-}2\%$, reflecting both analytical uncertainties and the spatial variation of Lu/Hf across many zircons; at the Lu/Hf ratios considered here, this contributes an uncertainty of $<0.1 \epsilon_{\text{Hf}}$ unit. For the calculation of ϵ_{Hf} values, we have adopted the chondritic values of Blichert-Toft et al.

(1997). To calculate model ages (T_{DM}) based on a depleted-mantle source, we have adopted a model with $^{176}\text{Hf}/^{177}\text{Hf}_i = 0.279718$ and $^{176}\text{Lu}/^{177}\text{Hf} = 0.0384$; this produces a value of $^{176}\text{Hf}/^{177}\text{Hf}$ (0.28325) similar to that of average MORB over 4.56 Ga. There are currently several values of the decay constant for ^{176}Lu : $1.93 \times 10^{-11}\text{yr}^{-1}$ proposed by Blichert-Toft et al., 1997; $1.865 \times 10^{-11}\text{yr}^{-1}$ by Scherer et al., 2001; and $1.983 \times 10^{-11}\text{yr}^{-1}$ by Bizzarro et al., 2003; calculations using these are provided in Table 3. Values used in Figure 4 have been calculated using the decay constant proposed by Scherer *et al.* (2001).

T_{DM} ages, which are calculated using the measured $^{176}\text{Lu}/^{177}\text{Hf}$ of the zircon, can only give a minimum age for the source material of the magma from which the zircon crystallised. Therefore we also have calculated, for each zircon, a “crustal” model age (T_{DM}^C , Table 3) which assumes that its parental magma was produced from an average continental crust ($^{176}\text{Lu}/^{177}\text{Hf} = 0.015$) that originally was derived from the depleted mantle.

References

- Bizzaro, M., Baker, J.A., Haack, H., Ulfbeck, D. & Rosing, M. Early History of Earth ‘s crust-mantle system inferred from hafnium isotope in chondrites. *Nature* 421, 931-933 (2003).
- Black L.P. & Gulson, B.L. The age of Mud Tank carbonatite, Strangways Range, North Territory. *BMR Journal of Australian Geol. and Gephys.* 3, 227-232 (1978).
- Blichert-Toft, J., Chauvel, C. & Albaréde, F. The Lu-Hf geochemistry of hondrites and the evolution of the mantle-crust system. *Earth and Planet. Sci. Letters*, 148, 243-258. Erratum. *Eart and Planet. Sci. Letters* 154, 349 (1997).
- Elhlou, S., Belousova, E.A., Griffin, W. L., Pearson, N. J. & O'Reilly, S.Y. Trace element and isotopic composition of GJ red zircon standard by Laser Ablation. Conference Abstract. *Geochim. Cosmochim. Acta* 70 (18), p. A158 (2006).
- Griffin, W.L. et al. The Hf isotope composition of cratonic mantle: LAM-MC-ICPMS analysis of zircon megacrysts in kimberlite. *Geochim. Cosmochim. Acta* 64, 133-147 (2000).
- Jackson, S. E., Pearson, N.J., Griffin, W.L. & Belousova, E.A. The application of laser ablation microprobe-inductively coupled plasma-mass spectrometry (LAM-ICP-MS) to in situ U-Pb zircon geochronology. *Chem. Geol.* 211, 47-69 (2004).
- Scherer, E., Münker, C. & Mezger, K., Calibration of the lutetium-hafnium clock. *Science* 293, 683-687 (2001).
- Wiedenbeck, M. et al. The natural zircon standards for U-Th-Pb, Lu-Hf, trace-element and REE analyses. *Geostands Newsletter* 19, 1-23 (1995).

Table DR1. Major and trace element composition of the studied rock types

Type of rock	eclogite/ ShS	Fe-Ti eclogite/UzS	garnetite/ UzS	plagiogranite/ UzS	TTG gneiss/ UzS	TTG gneiss/ ShS
Sample N	S-198/107	S-204/2b	S-204/23b	S-204/28	SB-810	KV-05/6
Age (Ga)	2.7	2.82	1.89	2.86	2.74*	2.96*
<i>Major elements (wt%)</i>						
SiO ₂	49.98	45.62	43.8	68.34	68.5	63.6
TiO ₂	0.76	2.22	1.34	0.49	0.41	0.63
Al ₂ O ₃	15.06	13.92	17.4	15.19	15.8	15.9
Fe ₂ O ₃	12.24	20.81	19.9	4.29	3.34	5.23
MnO	0.19	0.24	0.26	0.05	0.05	0.09
MgO	8.54	5.45	6.14	1.84	1.32	2.61
CaO	10.92	9.93	10.1	4.77	3.55	4.97
Na ₂ O	1.88	1.86	1.15	2.93	4.7	4.11
K ₂ O	0.31	0.03	0.07	0.06	1.76	2.17
P ₂ O ₅	0.07	0.05	<0.05	0.11	0.11	0.18
LOI	0.13	0.22	<1	0.52	0.28	0.35
Sum	100.08	100.00	100.00	100.15	99.82	99.84
Mg#	0.63	0.38	0.2	0.5	0.48	0.53
<i>Trace elements (ppm)</i>						
Sc	49	44	59.1	7	-	-
V	216	1170	512	54	38	46
Cr	478	6.5	24.2	37	8	12
Co	47.3	84	-	20	9	8.5
Ni	-	26	42.2	35	-	-
Rb	3.53	0.74	2.86	40	70	71
Sr	50.6	41	34.5	404	434	309
Y	16.6	28.9	25.9	8.9	6.87	7.4
Zr	43.4	77.1	39.5	227	146	147
Nb	2.46	3.75	1.84	6.72	6.1	5.71
Ba	25.7	5	22.5	536	627	529
La	1.41	0.76	0.31	25.4	30.8	22.2
Ce	4.49	2.31	0.35	50.5	57.6	43.2
Pr	0.71	0.44	0.08	5.54	6.46	4.53
Nd	3.84	3.22	0.52	20.2	21.5	15.9
Sm	1.34	2.16	0.7	3.6	3.18	2.81
Eu	0.44	1.04	0.49	1.29	0.85	0.75
Gd	1.63	3.29	2.41	3.34	2.54	2.55
Tb	0.36	0.76	0.66	0.45	0.34	0.34
Dy	2.64	4.95	4.23	2.04	1.5	1.67
Ho	0.63	1.07	0.91	0.34	0.27	0.29
Er	1.86	3.22	2.55	0.71	0.61	0.73
Tm	0.25	0.44	0.36	0.12	0.09	0.1
Yb	1.77	2.93	2.59	0.7	0.56	0.61
Lu	0.24	0.46	0.39	0.11	0.07	0.09
Hf	1.23	2.1	1.27	5.1	3.84	3.69
Ta	0.14	0.23	0.1	0.51	0.26	0.28
Th	0.08	0.13	<0.1	4.13	3.82	3.14
U	0.04	0.07	<0.1	1.14	0.29	0.28

* unpublished data of M. Mints et al., SHRIMP-II

Mg# = Mg/(Mg+Fe²⁺)

Table DR2A. Zircon LAM-ICPMS U-Pb isotopic data

Analysis #	$^{207}\text{Pb}/^{206}\text{Pb}$	$\pm 1\sigma$	$^{207}\text{Pb}/^{235}\text{U}$	$\pm 1\sigma$	$^{206}\text{Pb}/^{238}\text{U}$	$\pm 1\sigma$	$^{208}\text{Pb}/^{232}\text{Th}$	$\pm 1\sigma$	$^{207}\text{Pb}/^{206}\text{Pb}$	age	$^{207}\text{Pb}/^{235}\text{U}$	age	$^{206}\text{Pb}/^{238}\text{U}$	age	$^{208}\text{Pb}/^{232}\text{Th}$	age	$\pm 1\sigma$	Th(ppm)	U(ppm)	
<u>Sample: S198/107</u>																				
S198/107-01	0.1810	0.0020	12.72	0.12	0.5098	0.0042	0.1432	0.0025	2662	18	2659	9	2656	18	2704	44	110	828		
S198/107-02	0.1854	0.0021	13.23	0.13	0.5178	0.0047	0.1405	0.0028	2702	19	2697	9	2690	20	2658	49	53	353		
S198/107-03	0.1869	0.0021	13.50	0.13	0.5242	0.0047	0.1437	0.0027	2715	19	2715	9	2717	20	2714	47	68	362		
S198/107-05	0.1807	0.0021	12.69	0.14	0.5093	0.0049	0.1468	0.0032	2659	20	2657	10	2654	21	2769	57	76	617		
S198/107-06	0.1849	0.0020	13.24	0.12	0.5194	0.0045	0.1457	0.0023	2697	18	2697	9	2696	19	2750	40	57	417		
S198/107-07	0.1855	0.0020	13.35	0.13	0.5220	0.0046	0.1448	0.0023	2703	18	2705	9	2708	20	2733	41	75	513		
S198/107-09	0.1876	0.0024	13.57	0.18	0.5247	0.0057	0.1424	0.0039	2721	22	2720	12	2719	24	2691	69	47	146		
S198/107-10	0.1878	0.0023	13.72	0.15	0.5298	0.0046	0.1475	0.0037	2723	21	2730	10	2740	19	2781	64	28	67		
S198/107-11	0.1868	0.0023	13.39	0.15	0.5197	0.0049	0.1472	0.0036	2714	21	2707	11	2698	21	2776	63	55	377		
S198/107-14Core	0.1856	0.0022	13.31	0.14	0.5202	0.0048	0.1461	0.0034	2704	20	2702	10	2700	20	2756	60	183	814		
S198/107-14Rim	0.1846	0.0035	10.23	0.16	0.4020	0.0040	0.1105	0.0012	2695	32	2456	15	2178	18	2118	22	26	193		
S198/107-15Core	0.1801	0.0019	12.68	0.12	0.5105	0.0046	0.1439	0.0022	2654	18	2656	9	2659	20	2717	38	84	534		
S198/107-15Rim	0.1781	0.0026	11.95	0.13	0.4865	0.0048	0.1342	0.0014	2635	25	2600	10	2556	21	2545	25	120	665		
S198/107-16	0.1848	0.0021	12.86	0.13	0.5049	0.0047	0.1363	0.0026	2696	19	2670	10	2635	20	2583	47	201	911		
S198/107-17	0.1730	0.0020	11.39	0.13	0.4775	0.0049	0.1342	0.0029	2587	20	2556	11	2516	22	2546	52	93	731		
S198/107-18	0.1789	0.0022	12.10	0.14	0.4906	0.0051	0.1396	0.0032	2643	21	2612	11	2573	22	2641	56	39	103		
S198/107-19	0.1864	0.0021	13.36	0.13	0.5199	0.0045	0.1421	0.0028	2710	19	2705	9	2699	19	2686	50	56	376		
S198/107-20	0.1893	0.0022	13.84	0.15	0.5302	0.0050	0.1433	0.0029	2736	19	2739	10	2742	21	2707	51	76	431		
S198/107-21	0.1847	0.0021	13.21	0.13	0.5188	0.0046	0.1432	0.0027	2696	19	2695	9	2694	20	2705	48	116	730		
S198/107-22	0.1840	0.0021	13.19	0.12	0.5200	0.0044	0.1415	0.0026	2689	19	2694	9	2699	19	2674	47	119	456		
S198/107-23	0.1626	0.0021	9.90	0.12	0.4418	0.0047	0.1256	0.0030	2483	22	2426	12	2359	21	2392	53	59	101		
S198/107-24	0.1798	0.0024	12.49	0.17	0.5040	0.0057	0.1324	0.0038	2651	22	2642	13	2631	24	2513	68	90	495		
S198/107-25Core	0.1831	0.0021	13.27	0.15	0.5258	0.0053	0.1387	0.0029	2681	20	2699	11	2724	22	2626	52	82	247		
S198/107-25Rim	0.1843	0.0025	12.92	0.18	0.5088	0.0057	0.1433	0.0042	2692	23	2674	13	2651	24	2706	75	109	892		
S198/107-26	0.1861	0.0021	13.34	0.14	0.5200	0.0049	0.1455	0.0028	2708	19	2704	10	2699	21	2745	49	87	613		
S198/107-27	0.1813	0.0025	12.39	0.13	0.4958	0.0042	0.1365	0.0013	2665	23	2635	10	2596	18	2586	22	71	541		
S198/107-28	0.1833	0.0021	13.27	0.15	0.5250	0.0051	0.1516	0.0034	2683	20	2699	10	2720	22	2853	59	95	717		
S198/107-30	0.1407	0.0018	7.76	0.09	0.4003	0.0040	0.1156	0.0030	2235	22	2204	11	2170	19	2211	55	60	455		
S198/107-31	0.1846	0.0021	13.07	0.14	0.5135	0.0048	0.1420	0.0030	2694	19	2684	10	2671	20	2684	53	91	765		
<u>Sample: S204/2B</u>																				
S204/2B-1	0.1996	0.0028	14.97	0.22	0.5441	0.0065	0.1344	0.0042	2823	24	2813	14	2801	27	2549	74	247	178		
S204/2B-3	0.1684	0.0023	10.22	0.15	0.4401	0.0052	0.1265	0.0038	2542	24	2455	13	2351	23	2408	68	199	266		
S204/2B-4	0.1307	0.0054	7.07	0.28	0.3921	0.0050	0.1114	0.0013	2107	75	2120	35	2132	23	2135	24	472	366		
S204/2B-6	0.1303	0.0034	6.98	0.17	0.3886	0.0039	0.1105	0.0011	2102	47	2109	22	2116	18	2118	21	95	149		
S204/2B-7	0.1168	0.0013	5.53	0.06	0.3434	0.0033	0.1009	0.0019	1908	21	1905	9	1903	16	1942	34	177	224		
S204/2B-9	0.1984	0.0025	14.67	0.16	0.5361	0.0046	0.1503	0.0039	2813	21	2794	10	2767	19	2830	69	451	218		

S204/2B-11	0.1751	0.0025	11.96	0.17	0.4956	0.0058	0.1229	0.0040	2607	24	2602	14	2595	25	2344	71	242	268
S204/2B-12	0.1409	0.0037	7.67	0.18	0.3948	0.0043	0.1114	0.0012	2238	46	2193	21	2145	20	2134	22	335	345
S204/2B-16	0.1510	0.0057	8.14	0.29	0.3909	0.0045	0.1095	0.0012	2358	66	2247	33	2127	21	2101	22	246	242
S204/2B-17	0.1442	0.0019	7.88	0.10	0.3966	0.0042	0.1178	0.0032	2278	23	2218	12	2153	20	2251	57	107	198
S204/2B-19	0.1552	0.0031	9.23	0.16	0.4311	0.0042	0.1205	0.0012	2404	35	2361	16	2311	19	2299	22	119	207
S204/2B-23	0.1604	0.0019	9.18	0.10	0.4151	0.0040	0.1140	0.0023	2460	20	2356	10	2238	18	2182	41	82	183
S204/2B-24	0.1832	0.0026	12.09	0.17	0.4785	0.0055	0.1346	0.0044	2682	24	2611	13	2521	24	2552	77	486	522

Sample: S204/23B

S204/23B-1	0.1146	0.0022	5.12	0.09	0.3238	0.0040	0.0804	0.0062	1874	35	1839	16	1808	19	1563	116	1.2	12
S204/23B-2	0.1156	0.0024	5.44	0.11	0.3412	0.0044	0.0884	0.0042	1889	39	1891	17	1892	21	1712	78	3.2	11
S204/23B-5	0.1139	0.0017	5.28	0.08	0.3361	0.0037	2.4256	0.1724	1863	28	1865	12	1868	18	24887	1017	0.1	16
S204/23B-7	0.1152	0.0027	5.30	0.12	0.3336	0.0042	0.2072	0.0300	1884	43	1869	19	1856	20	3806	503	0.2	7
S204/23B-8	0.1150	0.0024	5.36	0.11	0.3380	0.0046	0.1693	0.0414	1880	39	1879	18	1877	22	3161	716	0.2	13
S204/23B-9	0.1157	0.0017	5.40	0.07	0.3385	0.0034	0.0934	0.0048	1890	26	1884	11	1879	16	1805	89	0.9	10
S204/23B-11	0.1156	0.0022	5.38	0.10	0.3377	0.0039	0.1168	0.0275	1889	36	1882	15	1875	19	2233	498	0.1	6
S204/23B-13	0.1154	0.0034	5.30	0.15	0.3331	0.0056	0.5185	0.2378	1887	55	1869	24	1853	27	8442	3165	0.0	8
S204/23B-14	0.1156	0.0023	5.41	0.10	0.3393	0.0043	0.0834	0.0065	1890	37	1886	16	1883	21	1619	122	1.0	9
S204/23B-15	0.1169	0.0021	5.60	0.09	0.3475	0.0040	0.0841	0.0124	1909	33	1916	15	1923	19	1633	230	0.3	6
S204/23B-16	0.1149	0.0022	5.28	0.10	0.3330	0.0042	0.1284	0.0141	1879	35	1865	16	1853	20	2441	252	0.5	14
S204/23B-17	0.1165	0.0020	5.44	0.09	0.3385	0.0039	0.0506	0.0640	1903	32	1891	14	1880	19	1050	1363	0.1	9
S204/23B-18	0.1160	0.0025	5.39	0.11	0.3372	0.0045	0.1262	0.0228	1896	39	1884	17	1873	22	2403	409	0.2	9
S204/23B-23	0.1168	0.0027	5.55	0.12	0.3448	0.0048	0.2191	0.2182	1907	42	1908	19	1910	23	4004	3617	0.02	8
S204/23B-25	0.1149	0.0033	4.75	0.13	0.3000	0.0049	0.0436	0.0030	1878	54	1776	23	1691	24	862	59	3.8	8

Sample: S204/28

S204/28-1	0.2038	0.0025	15.16	0.19	0.5397	0.0059	0.1437	0.0035	2856	20	2825	12	2782	25	2713	61	359	1079
S204/28-2	0.2035	0.0024	15.45	0.18	0.5507	0.0058	0.0674	0.0016	2855	20	2844	11	2828	24	1319	30	283	392
S204/28-3	0.1914	0.0021	12.37	0.13	0.4687	0.0046	0.1229	0.0024	2755	19	2633	10	2478	20	2342	43	168	585
S204/28-4	0.2060	0.0023	15.98	0.16	0.5628	0.0052	0.1408	0.0028	2874	18	2876	10	2878	22	2663	49	211	747
S204/28-5	0.2077	0.0025	16.23	0.19	0.5669	0.0058	0.1682	0.0039	2887	20	2891	11	2895	24	3143	67	131	183
S204/28-6	0.2042	0.0025	15.51	0.20	0.5509	0.0060	0.1491	0.0039	2860	21	2847	12	2829	25	2809	69	682	1825
S204/28-7	0.2005	0.0024	15.05	0.18	0.5444	0.0057	0.1365	0.0032	2830	20	2818	11	2802	24	2585	57	161	619
S204/28-8	0.2032	0.0026	16.01	0.21	0.5715	0.0063	0.1671	0.0048	2852	22	2878	13	2914	26	3123	84	628	818
S204/28-9	0.1943	0.0025	14.61	0.18	0.5453	0.0052	0.1512	0.0043	2779	22	2790	11	2806	22	2847	76	119	108
S204/28-11	0.2040	0.0026	15.77	0.20	0.5607	0.0059	0.1143	0.0033	2859	21	2863	12	2870	24	2187	59	84	211
S204/28-13	0.1826	0.0029	12.53	0.21	0.4977	0.0064	0.1429	0.0056	2676	27	2645	16	2604	28	2700	98	128	215
S204/28-14	0.2056	0.0032	16.06	0.26	0.5668	0.0069	0.1581	0.0063	2871	26	2881	16	2895	29	2967	110	400	1028
S204/28-16	0.1938	0.0031	13.85	0.23	0.5186	0.0066	0.1555	0.0062	2775	27	2740	16	2693	28	2922	108	77	244
S204/28-18	0.1946	0.0023	14.03	0.15	0.5230	0.0046	0.1469	0.0034	2781	20	2752	10	2712	20	2771	60	101	451
S204/28-19	0.2059	0.0024	15.74	0.18	0.5547	0.0056	0.1524	0.0032	2873	19	2861	11	2845	23	2867	57	272	925
S204/28-20	0.1950	0.0032	14.74	0.24	0.5481	0.0065	0.1680	0.0070	2785	28	2798	16	2817	27	3139	120	56	205
S204/28-22	0.1874	0.0038	12.64	0.21	0.4891	0.0056	0.1342	0.0017	2720	34	2653	16	2567	24	2546	30	129	535

S204/28-23	0.2073	0.0040	16.04	0.32	0.5617	0.0075	0.1581	0.0083	2884	32	2879	19	2874	31	2967	144	102	150
S204/28-24	0.2043	0.0027	15.38	0.20	0.5462	0.0060	0.1540	0.0045	2861	22	2839	13	2809	25	2895	79	64	196
S204/28-25	0.2039	0.0030	15.77	0.23	0.5610	0.0062	0.1598	0.0055	2857	24	2863	14	2871	26	2996	97	602	1606
S204/28-27Core	0.2101	0.0028	16.75	0.22	0.5783	0.0063	0.1251	0.0036	2906	22	2920	13	2942	26	2383	64	314	864
S204/28-27Rim	0.1996	0.0027	11.90	0.15	0.4326	0.0044	0.0769	0.0023	2823	22	2597	12	2318	20	1498	43	84	315
S204/28-28	0.2111	0.0027	16.67	0.21	0.5726	0.0061	0.1576	0.0043	2914	21	2916	12	2918	25	2958	75	104	345
S204/28-29	0.2036	0.0029	15.79	0.23	0.5626	0.0066	0.1581	0.0053	2856	24	2864	14	2877	27	2967	93	145	730
S204/28-31	0.2119	0.0034	14.81	0.23	0.5073	0.0058	0.1369	0.0052	2920	26	2803	15	2645	25	2593	93	150	181
S204/28-32Core	0.2064	0.0028	16.34	0.22	0.5744	0.0063	0.1249	0.0038	2877	23	2897	13	2926	26	2379	69	167	381
S204/28-33	0.2042	0.0047	15.60	0.35	0.5542	0.0078	0.1483	0.0094	2860	38	2853	21	2842	32	2794	165	59	102
S204/28-34	0.2028	0.0032	15.76	0.26	0.5638	0.0069	0.1605	0.0065	2849	27	2862	16	2882	29	3008	114	139	542

Table DR2B. Th-U-Pb isotopic data from SHRIMP-II (after Kaulina et al., 2007)

Analysis #	$^{206}\text{Pb}_c$ %	U ppm	Th ppm	$^{232}\text{Th}/^{238}\text{U}$	$^{206}\text{Pb}^*$ ppm	$^{206}\text{Pb}/^{238}\text{U}$ age $\pm 1\sigma$	$^{207}\text{Pb}/^{206}\text{Pb}$ (1) age $\pm 1\sigma$	Discordancy, %	$^{207}\text{Pb}/^{235}\text{U}$ (1)	$\pm\%$	$^{206}\text{Pb}^*/^{238}\text{U}$ (1)	$\pm\%$	err corr
------------	--------------------------	-------	--------	----------------------------------	----------------------------	--	---	-------------------	---	---------	---	---------	----------

Sample: S204/2B

Supposedly magmatic zircon (crystal with relics if magmatic zoning)

S204_2B.10.1	0,14	243	465	1,98	118	2891	± 50	2889	± 11	0	16.22	2.2	0.566	2.1	,951
Zircons recrystallized under eclogite facies conditions (light sections, Th/U > 1.75)															
S204_2B.3.1	0,15	496	857	1,79	211	2587	± 45	2787	± 8	8	13.28	2.2	0.494	2.1	,971
S204_2B.4.1	0,35	520	1488	2,95	234	2706	± 46	2801	± 11	3	14.16	2.2	0.522	2.1	,952
S204_2B.6.1	0,41	289	666	2,38	128	2665	± 47	2840	± 14	7	14.24	2.3	0.512	2.1	,932
S204_2B.7.1	0,22	432	971	2,33	186	2617	± 45	2799	± 9	7	13.58	2.2	0.501	2.1	,967
S204_2B.9.2	0,24	545	963	1,83	232	2588	± 45	2775	± 9	7	13.2	2.2	0.494	2.1	,965
S204_2B.10.2	0,23	383	920	2,48	163	2584	± 45	2802	± 9	8	13.39	2.2	0.493	2.1	,964
S204_2B.13.1	0,07	656	1458	2,30	267	2497	± 48	2750	± 7	10	12.45	2.4	0.473	2.3	,982

Zircons recrystallized strongly under eclogite facies conditions (light sections, Th/U < 1.75)

S204_2B.1.1	1,61	86	76	0,92	30.7	2210	± 44	2290	± 48	4	8.19	3.7	0.4089	2.4	,644
S204_2B.1.2	0,68	191	226	1,22	68.6	2234	± 41	2490	± 27	11	9.32	2.7	0.4141	2.2	,810
S204_2B.2.1	0,47	162	162	1,03	62.1	2361	± 44	2531	± 18	7	10.21	2.4	0.4423	2.2	,901
S204_2B.7.2	0,08	372	591	1,64	136	2280	± 43	2565	± 10	12	9.99	2.3	0.4243	2.2	,963
S204_2B.8.2	0,90	105	76	0,75	45.9	2622	± 50	2745	± 43	5	13.17	3.5	0.502	2.3	,663
S204_2B.8.3	0,30	560	563	1,04	213	2359	± 42	2553	± 13	8	10.33	2.3	0.4419	2.1	,934
S204_2B.9.1	0,76	189	142	0,78	72.3	2356	± 44	2561	± 28	9	10.37	2.8	0.4413	2.2	,794
S204_2B.11.1	0,37	299	321	1,11	120	2461	± 43	2586	± 13	5	11.08	2.3	0.4649	2.1	,939
S204_2B.15.1	0,07	309	368	1,23	130	2562	± 47	2675	± 11	4	12.28	2.3	0.488	2.2	,956

Thin light-gray rims of zircon crystals, Th/U < 0.5

S204_2B.2.2	2,67	34	7	0,22	11.1	2013	± 69	2081	± 160	3	6.51	9.9	0.366	4	,405
S204_2B.4.2	3,69	54	5	0,09	22.4	2437	± 62	2527	± 130	4	10.57	8.3	0.459	3	,366
S204_2B.5.1	1,18	79	23	0,30	30.2	2336	± 57	2504	± 70	7	9.92	5.1	0.437	2.9	,577
S204_2B.6.2	1,41	40	2	0,05	12.2	1949	± 49	2115	± 130	9	6.39	7.7	0.353	2.9	,379
S204_2B.12.1	1,42	29	8	0,27	11.7	2413	± 58	2576	± 64	7	10.76	4.8	0.454	2.9	,603
S204_2B.14.1	1,39	66	8	0,12	19.1	1839	± 43	2094	± 87	14	5.9	5.7	0.3302	2.7	,476

 Pb_c and Pb^* are common and radiogenic Pb, correspondingly,

Errors at standard calibration are from 0.37% to 0.91 %,

(1) correction for usual Pb was introduced using measured ^{204}Pb

Table DR3. Zircon Lu-Hf isotopic data

Analysis #	Measured Lu-Hf isotope ratios				Blichert-Toft et al. (1997) ^{176}Lu decay constant (1.93×10^{-11})				Scherer et al. (2001; 1.865×10^{-11})				Bizzarro et al. (2003; 1.983×10^{-11})				
	$^{176}\text{Hf}/^{177}\text{Hf}$	1 se	$^{176}\text{Lu}/^{177}\text{Hf}$	$^{176}\text{Yb}/^{177}\text{Hf}$	Hf _f	ε_{Hf}	1 se	T _{DM} (Ga)	T _{DM} ^C (Ga)	Hf _f	ε_{Hf}	T _{DM} (Ga)	T _{DM} ^C (Ga)	Hf _f	ε_{Hf}	T _{DM} (Ga)	T _{DM} ^C (Ga)
Sample: S198/107																	
S198/107-1	0.281117	0.000011	0.00013	0.004	0.281110	3.2	0.4	2.81	2.92	0.281111	1.0	2.91	3.09	0.281110	4.90	2.73	2.79
S198/107-2	0.281080	0.000009	0.00028	0.008	0.281065	2.5	0.3	2.87	2.99	0.281065	0.3	2.97	3.16	0.281064	4.26	2.79	2.86
S198/107-4	0.281075	0.000007	0.00026	0.008													
S198/107-5	0.281071	0.000009	0.00022	0.007	0.281059	1.3	0.3	2.88	3.03	0.281060	-0.9	2.98	3.20	0.281059	3.01	2.80	2.90
S198/107-6	0.281071	0.000007	0.00017	0.005	0.281062	2.3	0.2	2.87	3.00	0.281062	0.1	2.97	3.17	0.281062	4.05	2.80	2.87
S198/107-7	0.281068	0.000007	0.00028	0.007	0.281053	2.1	0.2	2.88	3.02	0.281053	-0.1	2.98	3.19	0.281052	3.86	2.81	2.88
S198/107-10	0.281127	0.000006	0.00029	0.009	0.281111	4.7	0.2	2.81	2.87	0.281112	2.5	2.91	3.04	0.281111	6.44	2.73	2.74
S198/107-11	0.281053	0.000008	0.00022	0.007	0.281041	1.9	0.3	2.90	3.03	0.281042	-0.2	3.00	3.21	0.281041	3.73	2.82	2.90
S198/107-12	0.281070	0.000010	0.00023	0.008													
S198/107-14	0.281080	0.000005	0.00021	0.007	0.281069	2.7	0.2	2.86	2.98	0.281069	0.5	2.96	3.15	0.281068	4.45	2.79	2.85
S198/107-15	0.281077	0.000006	0.00029	0.009	0.281062	1.2	0.2	2.87	3.03	0.281062	-0.9	2.97	3.20	0.281061	2.97	2.80	2.90
S198/107-16	0.281114	0.000008	0.00021	0.006	0.281103	3.7	0.3	2.82	2.91	0.281103	1.5	2.92	3.08	0.281102	5.46	2.74	2.78
S198/107-18	0.281114	0.000008	0.00021	0.006	0.281103	2.4	0.3	2.82	2.95	0.281103	0.3	2.92	3.12	0.281103	4.16	2.74	2.82
S198/107-20	0.281064	0.000009	0.00032	0.011	0.281047	2.7	0.3	2.89	3.01	0.281047	0.5	2.99	3.18	0.281046	4.46	2.81	2.87
S198/107-21	0.281052	0.000013	0.00016	0.004	0.281044	1.6	0.5	2.90	3.04	0.281044	-0.6	3.00	3.21	0.281043	3.36	2.82	2.91
S198/107-22	0.281151	0.000020	0.00018	0.006	0.281141	4.9	0.7	2.77	2.83	0.281142	2.7	2.87	3.00	0.281141	6.68	2.70	2.70
S198/107-23	0.281083	0.000017	0.00040	0.012	0.281063	-2.8	0.6	2.87	3.15	0.281064	-4.8	2.97	3.32	0.281063	-1.20	2.80	3.02
S198/107-24	0.281065	0.000009	0.00011	0.003	0.281059	1.1	0.3	2.88	3.04	0.281059	-1.1	2.98	3.21	0.281059	2.81	2.80	2.91
S198/107-25	0.281173	0.000015	0.00053	0.014	0.281145	4.8	0.5	2.77	2.83	0.281146	2.7	2.86	2.99	0.281144	6.58	2.69	2.70
S198/107-26	0.281058	0.000010	0.00021	0.006	0.281047	2.0	0.3	2.89	3.03	0.281047	-0.2	2.99	3.20	0.281047	3.78	2.81	2.89
S198/107-27	0.281055	0.000011	0.00021	0.006	0.281044	0.9	0.4	2.90	3.06	0.281044	-1.3	3.00	3.24	0.281043	2.60	2.82	2.93
S198/107-28	0.281086	0.000013	0.00013	0.004	0.281079	2.6	0.5	2.85	2.97	0.281080	0.4	2.95	3.14	0.281079	4.32	2.77	2.84
S198/107-30	0.281133	0.000014	0.00015	0.005	0.281126	-6.5	0.5	2.79	3.18	0.281127	-8.3	2.89	3.35	0.281126	-5.03	2.72	3.06
Sample: S204/2B																	
204/2B-1	0.281113	0.000012	0.00090	0.045	0.281063	5.3	0.4	2.87	2.91	0.281064	3.1	2.97	3.08	0.281061	7.15	2.79	2.78
204/2B-3	0.281143	0.000012	0.00059	0.027	0.281113	0.4	0.4	2.81	3.00	0.281114	-1.6	2.91	3.16	0.281113	2.03	2.73	2.87
204/2B-4	0.281143	0.000012	0.00059	0.027	0.281119	-9.8	0.4	2.81	3.29	0.281119	-11.4	2.91	3.45	0.281118	-8.44	2.73	3.16
204/2B-6	0.281281	0.000012	0.00184	0.086	0.281205	-6.8	0.4	2.72	3.10	0.281207	-8.4	2.81	3.25	0.281203	-5.56	2.65	2.99
204/2B-7	0.281100	0.000011	0.00076	0.039	0.281072	-16.2	0.4	2.88	3.52	0.281073	-17.6	2.98	3.68	0.281071	-14.95	2.80	3.39
204/2B-9	0.281226	0.000017	0.00151	0.081	0.281142	7.9	0.6	2.77	2.74	0.281145	5.7	2.86	2.90	0.281139	9.68	2.69	2.62
204/2B-11	0.281232	0.000016	0.00083	0.042	0.281189	4.6	0.6	2.71	2.78	0.281191	2.6	2.81	2.94	0.281188	6.31	2.64	2.66
204/2B-12	0.281288	0.000016	0.00121	0.045	0.281235	-2.6	0.6	2.66	2.94	0.281237	-4.3	2.76	3.10	0.281233	-1.14	2.59	2.83
204/2B-16	0.281144	0.000010	0.00124	0.048	0.281086	-5.0	0.4	2.86	3.19	0.281088	-6.8	2.96	3.35	0.281085	-3.49	2.78	3.06
204/2B-17	0.281144	0.000010	0.00124	0.048	0.281088	-6.8	0.4	2.86	3.24	0.281090	-8.6	2.96	3.40	0.281087	-5.37	2.78	3.11

204/2B-19	0.281235	0.000011	0.00171	0.064	0.281154	-1.5	0.4	2.77	3.01	0.281157	-3.3	2.87	3.16	0.281152	0.03	2.70	2.89
204/2B-23	0.281152	0.000013	0.00112	0.052	0.281093	3.0	0.5	2.84	2.94	0.281095	0.9	2.94	3.11	0.281091	4.72	2.76	2.82
204/2B-24	0.281289	0.000015	0.00108	0.041													

***Sample: S204/2B - calculated using 2820Ma crystallisation age**

204/2B-1	0.281113	0.000012	0.00090	0.045	0.281063	5.3	0.4	2.87	2.91	0.281065	3.0	2.97	3.08	0.281061	7.1	2.79	2.78
204/2B-3	0.281143	0.000012	0.00059	0.027	0.281110	6.9	0.4	2.81	2.80	0.281111	4.7	2.91	2.97	0.281109	8.8	2.73	2.68
204/2B-4	0.281143	0.000012	0.00059	0.027	0.281110	6.9	0.4	2.81	2.80	0.281111	4.7	2.91	2.97	0.281109	8.8	2.73	2.68
204/2B-6	0.281281	0.000012	0.00184	0.086	0.281178	9.4	0.4	2.72	2.65	0.281181	7.2	2.81	2.81	0.281175	11.1	2.65	2.53
204/2B-7	0.281100	0.000011	0.00076	0.039	0.281058	5.1	0.4	2.88	2.92	0.281059	2.8	2.98	3.09	0.281057	6.9	2.80	2.79
204/2B-9	0.281226	0.000017	0.00151	0.081	0.281142	8.1	0.6	2.77	2.73	0.281144	5.9	2.86	2.90	0.281139	9.8	2.69	2.61
204/2B-11	0.281232	0.000016	0.00083	0.042	0.281186	9.6	0.6	2.71	2.64	0.281187	7.4	2.81	2.80	0.281184	11.5	2.64	2.51
204/2B-12	0.281288	0.000016	0.00121	0.045	0.281221	10.9	0.6	2.66	2.56	0.281223	8.7	2.76	2.72	0.281219	12.7	2.59	2.44
204/2B-16	0.281144	0.000010	0.00124	0.048	0.281075	5.7	0.4	2.86	2.88	0.281077	3.5	2.96	3.05	0.281073	7.5	2.78	2.75
204/2B-17	0.281144	0.000010	0.00124	0.048	0.281075	5.7	0.4	2.86	2.88	0.281077	3.5	2.96	3.05	0.281073	7.5	2.78	2.75
204/2B-19	0.281235	0.000011	0.00171	0.064	0.281140	8.0	0.4	2.77	2.74	0.281143	5.8	2.87	2.90	0.281137	9.8	2.70	2.62
204/2B-23	0.281152	0.000013	0.00112	0.052	0.281090	6.2	0.5	2.84	2.85	0.281092	4.0	2.94	3.02	0.281088	8.0	2.76	2.72
204/2B-24	0.281289	0.000015	0.00108	0.041	0.281229	11.2	0.5	2.65	2.54	0.281231	9.0	2.75	2.70	0.281227	13.0	2.58	2.42

Sample: S204/23B

S-204/23B-1	0.281492	0.000014	0.00088	0.042	0.281460	-3.2	0.5	2.37	2.70	0.281461	-4.6	2.45	2.84	0.281459	-2.00	2.31	2.60
S-204/23B-2	0.281317	0.000013	0.00118	0.054	0.281273	-9.4	0.5	2.62	3.10	0.281275	-10.9	2.71	3.24	0.281272	-8.27	2.55	2.98
S-204/23B-5	0.281681	0.000015	0.00061	0.019	0.281659	3.6	0.5	2.11	2.27	0.281660	2.2	2.18	2.40	0.281658	4.82	2.05	2.18
S-204/23B-7	0.281911	0.000013	0.00001	0.000	0.281911	13.1	0.5	1.78	1.70	0.281911	11.6	1.84	1.81	0.281911	14.30	1.73	1.62
S-204/23B-8	0.281938	0.000011	0.00001	0.000	0.281938	13.9	0.4	1.74	1.65	0.281938	12.4	1.80	1.75	0.281938	15.16	1.69	1.57
S-204/23B-9	0.281329	0.000020	0.00101	0.048	0.281291	-8.8	0.7	2.60	3.06	0.281293	-10.2	2.69	3.20	0.281290	-7.60	2.53	2.94
S-204/23B-11	0.281900	0.000010	0.00003	0.001	0.281899	12.8	0.4	1.79	1.73	0.281899	11.3	1.85	1.83	0.281899	14.00	1.74	1.64
S-204/23B-13	0.281176	0.000014	0.00104	0.051	0.281137	-14.3	0.5	2.80	3.39	0.281139	-15.8	2.90	3.55	0.281136	-13.14	2.72	3.27
S-204/23B-15	0.281294	0.000016	0.00117	0.054	0.281250	-9.8	0.6	2.65	3.13	0.281252	-11.3	2.75	3.28	0.281249	-8.60	2.58	3.02
S-204/23B-16	0.281665	0.000017	0.00041	0.017	0.281650	3.7	0.6	2.12	2.28	0.281650	2.2	2.19	2.41	0.281649	4.89	2.06	2.19
S-204/23B-17	0.281441	0.000019	0.00160	0.080	0.281381	-5.3	0.7	2.49	2.85	0.281383	-6.7	2.57	2.99	0.281380	-4.11	2.42	2.75
S-204/23B-18	0.281877	0.000008	0.00000	0.000	0.281877	12.2	0.3	1.82	1.77	0.281877	10.6	1.88	1.88	0.281877	13.39	1.77	1.69
S-204/23B-23	0.281850	0.000010	0.00000	0.000	0.281850	11.5	0.4	1.86	1.82	0.281850	9.9	1.92	1.93	0.281850	12.70	1.81	1.74
S-204/23B-25	0.281251	0.000015	0.00122	0.052	0.281206	-12.1	0.5	2.71	3.25	0.281208	-13.5	2.81	3.40	0.281205	-10.92	2.64	3.13

Sample: S204/28

S204/28-1	0.281033	0.000010	0.00041	0.018	0.281010	4.2	0.4	2.94	3.00	0.281010	1.9	3.04	3.18	0.281009	6.10	2.86	2.87
S204/28-2	0.281023	0.000010	0.00054	0.019	0.280993	3.6	0.4	2.96	3.04	0.280994	1.3	3.06	3.21	0.280992	5.47	2.88	2.90
S204/28-3	0.281031	0.000015	0.00036	0.012	0.281011	1.9	0.5	2.94	3.07	0.281012	-0.3	3.04	3.24	0.281011	3.67	2.86	2.94
S204/28-4	0.281040	0.000009	0.00055	0.024	0.281009	4.6	0.3	2.94	2.99	0.281010	2.3	3.04	3.16	0.281008	6.51	2.86	2.85
S204/28-5	0.281049	0.000015	0.00039	0.016	0.281027	5.6	0.5	2.92	2.94	0.281027	3.3	3.02	3.11	0.281026	7.48	2.84	2.81
S204/28-6	0.281029	0.000017	0.00072	0.033	0.280988	3.6	0.6	2.97	3.04	0.280990	1.3	3.07	3.22	0.280987	5.43	2.89	2.91
S204/28-7	0.281035	0.000011	0.00074	0.035	0.280994	3.0	0.4	2.96	3.05	0.280995	0.8	3.06	3.23	0.280992	4.87	2.88	2.92

S204/28-8	0.281036	0.000016	0.00051	0.020	0.281007	4.1	0.6	2.94	3.01	0.281008	1.8	3.04	3.18	0.281006	5.92	2.86	2.87
S204/28-11	0.281014	0.000013	0.00042	0.017	0.280990	3.6	0.5	2.96	3.04	0.280991	1.3	3.07	3.22	0.280990	5.49	2.88	2.91
S204/28-13	0.281055	0.000025	0.00025	0.007	0.281042	1.0	0.9	2.90	3.06	0.281042	-1.1	3.00	3.23	0.281041	2.80	2.82	2.93
S204/28-14	0.281025	0.000013	0.00051	0.021	0.280996	4.1	0.5	2.96	3.02	0.280997	1.8	3.06	3.19	0.280995	6.00	2.88	2.88
S204/28-18	0.281045	0.000014	0.00038	0.017	0.281024	2.9	0.5	2.92	3.02	0.281025	0.7	3.02	3.20	0.281023	4.76	2.84	2.89
S204/28-19	0.281013	0.000014	0.00074	0.028	0.280971	3.3	0.5	2.99	3.07	0.280972	1.0	3.09	3.25	0.280970	5.13	2.91	2.94
S204/28-20	0.281141	0.000013	0.00067	0.024	0.281104	5.9	0.5	2.82	2.84	0.281105	3.7	2.92	3.01	0.281103	7.69	2.74	2.72
S204/28-23	0.281051	0.000014	0.00041	0.012	0.281028	5.6	0.5	2.92	2.94	0.281029	3.2	3.02	3.11	0.281027	7.45	2.84	2.81
S204/28-25	0.281061	0.000011	0.00052	0.022	0.281031	5.0	0.4	2.91	2.95	0.281032	2.8	3.01	3.12	0.281031	6.90	2.83	2.82
S204/28-31	0.281044	0.000025	0.00027	0.009	0.281028	6.4	0.9	2.91	2.91	0.281029	4.1	3.02	3.09	0.281028	8.37	2.84	2.78
S204/28-34	0.281059	0.000024	0.00075	0.025	0.281017	4.3	0.8	2.93	2.99	0.281018	2.1	3.03	3.16	0.281015	6.16	2.85	2.86

Table DR4. Calculated P-T conditions for the inferred equilibria in the studied samples (also see Figure 4)

Lithology, sample	Stage, assemblage	Calibration	T (°C)	P (GPa)
<i>Shirokaya Salma locality</i>				
Eclogite	<i>Retrograde</i>			
S-198/107	Grt-Cpx-Pl-Qtz	GCPQ (TPF)	723	1.15
	Cpx-Pl-Qtz	McCarthy, Patiño Douce, equation 5, (1998)	(750)	1.06
	Cpx-Pl-Qtz	McCarthy, Patiño Douce, equation 7, (1998)	(750)	0.97
<i>Uzkaya Salma locality</i>				
Eclogite	<i>Peak</i>			
SB-812	Cpx-Pl-Qtz	Holland (1980)	(700)	1.28
	Grt-Cpx	Powell (1985)	698	(1.30)
	<i>Retrograde</i>			
	Cpx-Pl-Qtz	McCarthy, Patiño Douce, equation 5, (1998)	(750)	1.03
	Cpx-Pl-Qtz	McCarthy, Patiño Douce, equation 7, (1998)	(750)	0.95
Eclogite, M2a	<i>Prograde (?)</i>			
	Included in garnet			
	Grt-Hbl	Powell (1985)	687	
	Grt-Hbl	Lavrentieva and Perchuk (1989)	704	
	Grt-Cpx	Powell (1985)	707	(1.0)
	<i>Peak</i>			
	Cpx-Pl-Qtz	Holland (1980)	(750)	(1.4)
	<i>Retrograde</i>			
	Grt-Cpx-Pl-Qtz	GCPQ (TPF)	747	1.19
	Cpx-Pl-Qtz	McCarthy, Patiño Douce, equation 5, (1998)	(750)	1.32
	Cpx-Pl-Qtz	McCarthy, Patiño Douce, equation 7, (1998)	(750)	1.2
Eclogite, S204/16	<i>Prograde (?)</i>			
	Included in garnet			
	Grt-Hbl	Powell (1985)	643	
	Grt-Hbl	Lavrentieva and Perchuk (1989)	670	
	Hbl-Pl-Qtz	Holland and Blundy (1994)	650	(0.7)
	Hbl-Pl	Holland and Blundy (1994)	490	(0.7)
	Cpx-Pl-Qtz	McCarthy, Patiño Douce, equation 5, (1998)	(700)	0.74
	Cpx-Pl-Qtz	McCarthy, Patiño Douce, equation 7, (1998)	(700)	0.66
Fe-Ti eclogite, S204/3	<i>Retrograde</i>			
	Grt-Cpx-Pl-Qtz	GCPQ (TPF)	689	1.0
	Cpx-Pl-Qtz	McCarthy, Patiño Douce, equation 5, (1998)	(700)	(0.85)
	Cpx-Pl-Qtz	McCarthy, Patiño Douce, equation 7, (1998)	(700)	(0.75)

Ky-Grt-Bt plagiogranite		<i>Peak</i>		
S204/28	Grt-Bt	GB (TPF)	732	(1.4)
	Grt-Ky-Pl-Qtz	Koziol and Newton (1989)	(700)	1.35
<i>Retrograde</i>				
	Grt-Bt	GB (TPF)	643	(1.0)
	Grt-Ky-Pl-Qtz	Koziol and Newton (1989)	600	0.91

Abbreviations for the self-consistent calibrations of the program TPF (Fonarev et al., 1991; Konilov, 1999; Maaskant, 2004): GCPQ is Grt-Cpx-Pl-Qtz geothermobarometer combined of Grt-Cpx-Pl-Qtz geobarometer (Fonarev et al., 1994) and Grt-Cpx geothermometer (Powell, 1985); GB is Grt-Bt geothermometer, averaged of (Perchuk and Lavrent'eva, 1983) and (Holdaway and Lee, 1977).

References to Table DR4:

- Fonarev, V.I., Graphchikov, A.A., and Konilov, A.N., 1991. A consistent system of geothermometers for metamorphic complexes. International Geology Review 33, 8, 743-783.
- Fonarev, V.I., Graphchikov, A.A., and Konilov A.N., 1994. Experimental studies of equilibria with minerals of variable composition and geological thermobarometry. Experimental problems of geology. Moscow, Nauka. 323-355 (in Russian).
- Holdaway, M.J., and Lee, S.M., 1977. Fe-Mg cordierite stability in high-grade pelitic rocks based on experimental, theoretical, and natural observations. Contrib. Mineral. Petrol. 63, 175-198.
- Holland, T.J.B., 1980. The reaction albite=jadeite+quartz determined experimentally in the range 600–1200°C. Am. Min. 65, 129–134.
- Holland, T., and Blundy J., 1994. Non-ideal interactions in calcic amphiboles and their bearing on amphibole-plagioclase thermometry. Contrib. Miner. Petrol. 116, 433-447.
- Konilov, A.N., 1999. Testing of the consistent system of geothermometers and geobarometers of the program TPF. Experiment in GeoSciences 8, 1, 60-62.
- Koziol, A.M., and Newton, R.C., 1989. Grossular activity-composition relationships in ternary garnets determined by reversed displaced-equilibrium experiments. Contrib. Miner. Petrol. 103, 423-433.

Lavrent'eva, I.V. and Perchuk, L.L., 1989. An experimental study of the amphibole-garnet equilibrium: calcium free system Dokl. Akad. Nauk USSR 306, 173-175 (in Russian).

Maaskant, P., 2004. Thermobarometry of the Furua granulites, Tanzania: a comparative study. Neue Jb. Miner. Abh., Stuttgart, 180, 65-100.

McCarthy, T.C., and Patiño Douce, A.E., 1998. Empirical calibration of the silica-Ca-tschermark's-anorthite (SCAn geobarometer). J. Metamorph. Geology 6, 671-682.

Perchuk, L.L., and Lavrent'eva, I.V., 1983. Experimental investigation of exchange equilibria in the system cordierite-garnet-biotite. Advances in Physical Geochemistry 3, 199-239.

Powell, R., 1985. Regression diagnostic and robust regression in geothermometer/geobarometer calibration: the garnet-clinopyroxene geothermometer revised. J. Metamorph. Geology 3, 231-243.




Figure DR1. Regional tectonic map of the eastern Fennoscandian Shield (left) and geologic sketch-map of the area of the Archean eclogite localities (right), where

Meso-Neoarchean:

- (1) – Keret' tectonic nappe at southern margin of the Archean Kola continent, 2.89-2.70 Ga TTG;
- (2) – ≥ 2.88 -2.85 Ga Central Belomorian suture zone;

- (3) – eclogite localities (UzS – Uzkaya Salma and ShS – Shirokaya Salma);
- (4) – 2.78-2.74 Ga island arc type greenstones;
- (5-7) – Belomorian orogen: 3.1-2.77 Ga TTG (5), 2.80-2.77 Ga greenstones (6), and 2.78 Ga Iringora ophiolites (see Shchipansky et al., 2004) (7);
- (8) – Karelia continent, 3.5-2.7 Ga TTG.

Neoarchean:

- (9) – 2.74-2.68 Ga Chupa granulite-gneiss belt;
- (10) – 2.73-2.71 Ga enderbite bodies;
- (11) – 2.72-2.66 Ga greenstones.

Paleoproterozoic:

- (12-13) – 2.53-2.42 Ga intrusive bodies: layered mafic-ultramafics (12) and gabbro-anorthosites (13);
- (14) – ca. 1.95 Ga Lapland-Kolvitsa granulite-gneiss belt;
- (15) – Paleoproterozoic volcano-sedimentary belts;
- (16) – late Paleoproterozoic granite-gneiss domes;
- (17-19) – faults: overthrusts (17), normal faults and strike-slip faults (18), faults beneath water areas (19).




Figure DR2. Geological and petrological features of the Salma eclogites. a) A photo of the UzS body in outcrop; b) almandine-pyrope garnets set in a matrix of vermicular clinopyroxene-plagioclase symplectite and locally preserved omphacite; backscattered electron image (left) and transmitted light image (right).

Inclusions of the relic low-T minerals in the Salma eclogite as an evidence of the prograde metamorphic evolution: c) atoll type garnet within Cpx-Pl symplectite microphoto (left) with the inset showing enlarged BSE image of the central part of atoll, and a compositional profile (right), where shaded area corresponds to “lagoon” of atoll; d) BSE image of the atoll type garnet (left) with inclusions marked by white arrows for Ab (albite) and Pmp (pumpellyite), and a compositional profile (right), where shaded area corresponds to a zone with relics of Ab, Act (actinolite) and Pmp, scale bar is 2 mm long; e) SE image showing quartz needle-shaped inclusions (rods) in omphacite, width of a field of view is 200 μm ; f) photomicrograph of the quartz needle-shaped inclusions in omphacite, scale bar is 50 μm long. Similar inclusions are

known in the UHP assemblages, e.g., in Kokchetav massif (Katayama, Maruyama, 2009), Pohorje, Eastern Alps (Janák et al., 2004), Alpe Arami (Dobrzhinetskaya et al., 2002), Blumenau eclogite, Erzgebirge (Chopin, Ferraris, 2003). This feature permits suggestion that peak conditions in the history of the Salma eclogite could reach much higher pressure than 1.4 GPa that were fixed in these rocks to-day. Also see Table DR4 and Fig. 4.

References in Data Repository Figure DR2:

- Chopin, C., Ferraris, G. Mineral chemistry and mineral reactions in UHPM rocks. In: (Carswell D.A., Compagnoni R., Rolfo F., Eds.) Ultrahigh Pressure Metamorphism. EMU Notes in Mineralogy, Vol. 5 (2003), Chapter 7, 191–227.
- Dobrzhinetskaya, L.F., Schweinehage, R., Massonne, H.-J., Green, H.W. Silica precipitates in omphacite from eclogite at Alpe Arami, Switzerland: evidence of deep subduction. *J. metamorphic Geol.*, 2002, 20, 481–492.
- Janák, M., Froitzheim, N., Lupták, B., Vrabec, M., Krogh Ravna, E.J. First evidence for ultrahigh-pressure metamorphism of eclogites in Pohorje, Slovenia: Tracing deep continental subduction in the Eastern Alps. *Tectonics*, 2004, v.23, TC5014, doi: 10.1029/2004TC001641.
- Katayama, I., Maruyama, S. Inclusion study in zircon from ultrahigh-pressure metamorphic rocks in the Kokchetav massif: an excellent tracer of metamorphic history. *Journal of the Geological Society, London*, 2009, v.166, no.4, 783–796, doi: 10.1144/0016-76492008-019.
- Shchipansky, A.A., Samsonov, A.V., Bibikova, E.V., Babarina, I.I., Konilov, A.N., Krylov, K.K., Slabunov, A.I., Bogina, M.M. 2.8 Ga boninite-hosting partial suprasubduction zone ophiolite sequences from the North Karelian greenstone belt, NE Baltic Shield, Russia. In: Precambrian ophiolites and related rocks (T. Kusky – Ed.). 2004. Development in Precambrian Geology, 13. Elsevier, Amsterdam, Chapter 14, 425-486.




Figure DR3

Figure DR3. Backscattered electron/cathodoluminescence images of representative zircon grains from the analysed samples. Scale bar: 100 µm for images a, c, d, e and f and 50 µm for images b, g and h. See text for explanations.