

Paleomagnetic Data Analysis of Late Stage Volcanics from the Midcontinent Rift

This Jupyter notebook is provided as the main Data Repository for a manuscript in review entitled **The end of Midcontinent Rift magmatism and the paleogeography of Laurentia** by Luke M. Fairchild, Nicholas L. Swanson-Hysell, Jahandar Ramenzani, Courtney J. Sprain, and Samuel A. Bowring. This notebook can be [downloaded for interactive viewing¹](#) or [viewed statically within a web browser²](#); it is highly recommended that this notebook be viewed in either of these two ways, although a PDF rendering of the notebook is also provided in the Data Repository.

¹https://github.com/Swanson-Hysell-Group/2016_Late_Rift/blob/master/Code/Late_Rift_Data_Analysis.ipynb

²https://nbviewer.jupyter.org/github/swanson-hysell-group/2016_Late_Rift/blob/master/Code/Late_Rift_Data_Analysis.ipynb

Contents

1 Import and develop functions for use within the Jupyter notebook	2
2 Schroeder-Lutsen Basalts	2
2.1 Prior data	2
2.2 New data from Two Island River	6
2.2.1 Divergent VGP populations from new SLB data	12
2.3 Combine new SLB data with Tauxe and Kodama (2009)	20
3 Lake Shore Traps	23
4 Michipicoten Island	28
4.1 Palmer and Davis (1987) data	28
4.2 New data from the Michipicoten Island Formation	31
4.3 Combine new Michipicoten data with Palmer and Davis (1987)	38
5 Paleogeography of Laurentia	42
5.1 Late Rift Poles Compilation	42
5.2 Rates of plate motion	46
5.2.1 Latitudinal motion implied by Osler and Michipicoten poles	46
5.2.2 Latitudinal motion implied by Mamaing lowerN/upperR and Michipicoten poles	47
5.2.3 Latitudinal motion implied by North Shore Volcanic Group and Michipicoten poles	48
5.3 Age of Keweenawan sediments	49
5.3.1 <i>Nonesuch Shale</i>	49
5.3.2 <i>Jacobsburg Sandstone</i>	51
6 Map of late stage volcanics paleomagnetic sites	57
7 References	59

1 Import and develop functions for use within the Jupyter notebook

The code block below imports necessary libraries and defines functions that will be used in the data analysis below. The modules `pmag.py`, `pmagplotlib.py` and `ipmag.py` are from the PmagPy software package (<https://github.com/ltauxe/PmagPy>; Tauxe et al., 2016). The current version of these modules (PmagPy 3.4) are included in the repository with this notebook. Other necessary function libraries (`matplotlib`, `pandas`, `Basemap`, `IPython`, `numpy`, `scipy`) are included with standard scientific Python distributions.

```
In [1]: import pmagpy_3_4.pmag as pmag
import pmagpy_3_4.pmagplotlib as pmagplotlib
import pmagpy_3_4.ipmag as ipmag

import matplotlib.pyplot as plt
from mpl_toolkits.basemap import Basemap
from matplotlib.patches import Polygon
import matplotlib.patches as mpatch
from matplotlib.collections import PatchCollection
import pandas as pd
from IPython.core.display import HTML
import numpy as np
import scipy as sp
from scipy import special
from IPython.display import Image
```

This notebook runs with figures inline in the Jupyter notebook (instead of opening up in another window) by executing this command:

```
In [2]: %matplotlib inline
%config InlineBackend.figure_formats = {'svg',}
```

Throughout this analysis there will be maps that show pole positions. The `pole_figure_appearance` function controls aspects of the appearance of these maps and eliminates the need to make these customizations for every plot.

```
In [3]: def pole_figure_appearance(size = (7,7)):
    m = Basemap(projection='ortho',lat_0=35,lon_0=200,resolution='c',
                area_thresh=50000)
    plt.figure(figsize=size)
    m.drawcoastlines(linewidth=0.25)
    m.fillcontinents(color='bisque',lake_color='white',zorder=1)
    m.drawmapboundary(fill_color='white')
    m.drawmeridians(np.arange(0,360,30))
    m.drawparallels(np.arange(-90,90,30))
    return m
```

2 Schroeder-Lutsen Basalts

2.1 Prior data

Tauxe and Kodama (2009) published data from sites of the North Shore Volcanic Group and Schroeder-Lutsen Basalts that was the first study of these units to use modern methods. Books (1968, 1972) and Palmer (1970) also developed data that can largely be considered to be superseded by the Tauxe and Kodama results. In the code below, the (Tauxe and Kodama, 2009) data are imported and then the data that are from the Schroeder-Lutsen Basalts are split off from the rest of the data that were generated by (Tauxe and Kodama, 2009) from the North Shore Volcanic Group.

```
In [4]: Tauxe_NSVG_Data=pd.read_csv('../Data/Previous_studies/Tauxe2009a_data.csv',sep=',')
Tauxe_NSVG_Data.head()
```

Out[4]: MagIC contribution of Tauxe and Kodama (2009) (<http://earthref.org/MagIC/8407/>)

NOTE: The hyperlink above may only be active if you are reading this in Adobe Acrobat. If you are having trouble opening the local file via this link, or if you have downloaded this document separately from the accompanying data files, the file can be found in the Data Repository under /Data/Previous_studies/Tauxe2009a_data.csv and opened manually.

The **sequence** column contains an assignment of the sequence with 'nsl' signifying that we interpret the flow to be within the Schroeder-Lutsen basalts. A new dataframe can be made with only these flows.

```
In [5]: NSVG_nswu = Tauxe_NSVG_Data.ix[Tauxe_NSVG_Data['sequence'] == 'nswu']
NSVG_nswu.reset_index(inplace=True)

NSVG_nswu_VGPs = ipmag.make_di_block(NSVG_nswu['pole_lon'],NSVG_nswu['pole_lat'])
NSVG_nswu_mean = pmag.fisher_mean(NSVG_nswu_VGPs)
ipmag.print_pole_mean(NSVG_nswu_mean)


NSVG_nsl = Tauxe_NSVG_Data.ix[Tauxe_NSVG_Data['sequence'] == 'nsl']
NSVG_nsl.reset_index(inplace=True)

NSVG_nsl_VGPs = ipmag.make_di_block(NSVG_nsl['pole_lon'],NSVG_nsl['pole_lat'])
NSVG_nsl_mean = pmag.fisher_mean(NSVG_nsl_VGPs)
ipmag.print_pole_mean(NSVG_nsl_mean)

m = pole_figure_appearance()



ipmag.plot_vgp(m,NSVG_nswu['pole_lon'].tolist(),NSVG_nswu['pole_lat'].tolist(),
                color='b',label='NSVG (nswu; Tauxe and Kodama (2009))')
ipmag.plot_pole(m,NSVG_nswu_mean['dec'],NSVG_nswu_mean['inc'],
                NSVG_nswu_mean['alpha95'], marker='s',color='b')
ipmag.plot_vgp(m,NSVG_nsl['pole_lon'].tolist(),NSVG_nsl['pole_lat'].tolist(),
                color='g',label='SLB (nsl; Tauxe and Kodama (2009))')
ipmag.plot_pole(m,NSVG_nsl_mean['dec'],NSVG_nsl_mean['inc'],
                NSVG_nsl_mean['alpha95'], marker='s',color='g')
plt.legend(loc='upper center')
plt.show()
```

Plong: 182.1 Plat: 35.8
Number of directions in mean (n): 47
Angular radius of 95% confidence (A_95): 3.1
Precision parameter (k) estimate: 45.7
Plong: 180.5 Plat: 28.8
Number of directions in mean (n): 10
Angular radius of 95% confidence (A_95): 6.5
Precision parameter (k) estimate: 55.6



```
In [6]: ipmag.fishqq(NSVG_ns1['pole_lon'].tolist(),NSVG_ns1['pole_lat'].tolist())
```

```
Out[6]: {'Dec': 180.5436552009686,
 'Inc': 28.803422228130206,
 'Me': 0.65209594384149339,
 'Me_critical': 1.094,
 'Mode': 'Mode 1',
 'Mu': 0.98495553684353243,
 'Mu_critical': 1.207,
 'N': 10,
 'Test_result': 'consistent with Fisherian model'}
```



2.2 New data from Two Island River

We present new paleomagnetic data from 40 Schroeder-Lutsen basalt flows exposed within Two Island River. Magnetizations held by magnetite and those held by hematite or maghemite are virtually identical (see equal area plots below), which suggests oxidation of these lava flows in association with eruption or shortly thereafter.

```
In [7]: SLB_Data_all = pd.read_csv('../Data/SLB/pmag_results.txt',sep='\t',skiprows=1)
    SLB_Data_all_tc = SLB_Data_all.ix[SLB_Data_all['tilt_correction'] == 100.0]
    SLB_Data_hem = SLB_Data_all.ix[SLB_Data_all['pole_comp_name'] == 'hem']
    SLB_Data_hem = SLB_Data_hem.ix[SLB_Data_hem['tilt_correction'] == 100.0]
    SLB_Data_hem.reset_index()
    SLB_Data = SLB_Data_all.ix[SLB_Data_all['pole_comp_name'] == 'mag']
    SLB_Data = SLB_Data.ix[SLB_Data['tilt_correction'] == 100.0]
    SLB_Data.reset_index()
    SLB_Data.head()
```

```
Out[7]: View raw data in the MagIC contribution of this study
        (https://earthref.org/MagIC/doi/10.1130/L580.1)
```

If you are having trouble opening the file via this link, the file can be found in the Data Repository under /Data/SLB/pmag_results.

```
In [8]: plt.figure(num=1,figsize=(6,6))
    ipmag.plot_net(1)

    SLB_dec = SLB_Data['average_dec'].tolist()
    SLB_inc = SLB_Data['average_inc'].tolist()
    SLB_a95 = SLB_Data['average_alpha95'].tolist()


    for i in range(len(SLB_dec)):
        ipmag.plot_di_mean(SLB_dec[i], SLB_inc[i], SLB_a95[i])

    SLB_mean_dir = ipmag.fisher_mean(SLB_dec,SLB_inc)

    ipmag.plot_di_mean(SLB_mean_dir['dec'],SLB_mean_dir['inc'],
                       SLB_mean_dir['alpha95'],marker='s', color='r')
    plt.title('Schroeder-Lutsen magnetite directions')

    plt.savefig('Code_output/All_SLB_mag_data.pdf')
    plt.show()
```

Schroeder-Lutsen magnetite directions



```
In [9]: plt.figure(num=1,figsize=(6,6))
ipmag.plot_net(1)

SLB_hem_dec = SLB_Data_hem['average_dec'].tolist()
SLB_hem_inc = SLB_Data_hem['average_inc'].tolist()
SLB_hem_a95 = SLB_Data_hem['average_alpha95'].tolist()


for i in range(len(SLB_hem_dec)):
    ipmag.plot_di_mean(SLB_hem_dec[i], SLB_hem_inc[i], SLB_hem_a95[i])

SLB_hem_mean_dir = ipmag.fisher_mean(SLB_hem_dec,SLB_hem_inc)

ipmag.plot_di_mean(SLB_hem_mean_dir['dec'],SLB_hem_mean_dir['inc'],
                    SLB_hem_mean_dir['alpha95'],marker='s', color='r')
```

```
plt.title('Schroeder-Lutsen hematite/maghemite directions')  
plt.savefig('Code_output/All_SLB_hem_data.pdf')  
plt.show()
```

Schroeder-Lutsen hematite/maghemite directions



Below, we calculate the average angular difference between the magnetite and hematite paleomagnetic directions of each sampled flow.

```
In [10]: SLB_samples = pd.read_csv('../Data/SLB/pmag_specimens.txt', sep='\t', skiprows=1)  
SLB_samples = SLB_samples[SLB_samples['specimen_tilt_correction'] == 100.0]  
  
In [11]: SLB_mag_hem_diff = []  
for sample in SLB_samples['er_specimen_name'].unique().tolist():  
    try:  
        mag_dec = float(SLB_samples.loc[SLB_samples['er_specimen_name'] == sample] \
```

```
        .loc[SLB_samples['specimen_comp_name']=='mag'].specimen_dec)
mag_inc = float(SLB_samples.loc[SLB_samples['er_specimen_name']==sample]\n                  .loc[SLB_samples['specimen_comp_name']=='mag'].specimen_inc)
hem_dec = float(SLB_samples.loc[SLB_samples['er_specimen_name']==sample]\n                  .loc[SLB_samples['specimen_comp_name']=='hem'].specimen_dec)
hem_inc = float(SLB_samples.loc[SLB_samples['er_specimen_name']==sample]\n                  .loc[SLB_samples['specimen_comp_name']=='hem'].specimen_inc)
SLB_mag_hem_diff.append(pmag.angle([mag_dec, mag_inc], [hem_dec, hem_inc]))
except:
    continue
```


Average angle between magnetite and hematite/maghemite fits for all samples: 2.86292160508

```
In [12]: SLB_Data_low_lat = SLB_Data.ix[SLB_Data['vgp_lat'] < 15]
          SLB_Data_hi_lat = SLB_Data.ix[SLB_Data['vgp_lat'] >= 15]

new_SLB_mean = ipmag.fisher_mean(SLB_Data['vgp_lon'].tolist(),
                                  SLB_Data['vgp_lat'].tolist())
new_SLB_low_lat_mean = ipmag.fisher_mean(SLB_Data_low_lat['vgp_lon'].tolist(),
                                         SLB_Data_low_lat['vgp_lat'].tolist())
new_SLB_hi_lat_mean = ipmag.fisher_mean(SLB_Data_hi_lat['vgp_lon'].tolist(),
                                         SLB_Data_hi_lat['vgp_lat'].tolist())

m = pole_figure_appearance()

# ipmag.plot_vgp(m,SLB_Data['vgp_lon'].tolist(),SLB_Data['vgp_lat'].tolist(),
#                 label='SLB VGPs',color='r')
ipmag.plot_vgp(m,SLB_Data_low_lat['vgp_lon'].tolist(),
                SLB_Data_low_lat['vgp_lat'].tolist(),
                label='SLB low lat VGPs',color='b')
ipmag.plot_pole(m,new_SLB_low_lat_mean['dec'],new_SLB_low_lat_mean['inc'],
                 new_SLB_low_lat_mean['alpha95'],
                 label='SLB low lat mean',marker='s',color='b')
ipmag.plot_vgp(m,SLB_Data_hi_lat['vgp_lon'].tolist(),
                SLB_Data_hi_lat['vgp_lat'].tolist(),
                label='SLB high lat VGPs',color='g')
ipmag.plot_pole(m,new_SLB_hi_lat_mean['dec'],new_SLB_hi_lat_mean['inc'],
                 new_SLB_hi_lat_mean['alpha95'],
                 label='SLB high lat mean',marker='s',color='g')
ipmag.plot_pole(m,new_SLB_mean['dec'],new_SLB_mean['inc'],new_SLB_mean['alpha95'],
                 label='All SLB mean',marker='s',color='red')
```



As is apparent in the plots above, SLB paleomagnetic directions fall into two distinct populations. We organize these into lower and higher latitude poles below.

In [13]: `SLB_Data_low_lat.head()`

Out[13]: VGP Site: SLB01
VGP Site: SLB02
VGP Site: SLB10
VGP Site: SLB11
VGP Site: SLB12


In [14]: `SLB_Data_hi_lat.head()`


Out[14]: VGP Site: SLB03
VGP Site: SLB04
VGP Site: SLB05
VGP Site: SLB06
VGP Site: SLB07

As seen below, a Fisherian distribution is rejected for our new Schroeder-Lutsen data (Fisher et al., 1987).

```
In [15]: # Test whether new SLB data qualifies as a Fisherian distribution
ipmag.fishqq(SLB_Data['vgp_lon'].tolist(),SLB_Data['vgp_lat'].tolist())
```

```
Out[15]: {'Dec': 189.55557939703078,
'Inc': 26.745255032788066,
'Me': 0.53371584170306374,
'Me_critical': 1.094,
'Mode': 'Mode 1',
'Mu': 1.7737870824702469,
'Mu_critical': 1.207,
'N': 40,
'Test_result': 'Fisherian model rejected'}
```






When the two populations are grouped, the combined SLB data fail the Fisher quantile-quantile test above. Possible explanations for this non-Fisherian distribution are explored below.

2.2.1 Divergent VGP populations from new SLB data

Rapid multiple-eruption episodes that are causing excursion behavior to be overrepresented in the dataset This explanation would likely predict that the more southerly VGPs are excursion and therefore stratigraphically grouped. However, approaching the problem from a stratigraphic standpoint reveals that the low latitude VGPs are encountered in multiple parts of the stratigraphic succession. Since these VGPs are coming from flows throughout SLB stratigraphy (some sequential flows, others isolated) if the southerly population is excursion it would have needed to be a direction that was returned to repeatedly by the field. In the stratigraphic column below, the blue arrows mark sampled flows that exhibit low latitude directions, and the red arrows mark sampled flows that exhibit high latitude directions. It is quite clear from this figure that there is no stratigraphic pattern of the anomalous paleomagnetic directions.



Additionally, a broader map view of the stack of lava flows highlights the discontinuity of the two VGP populations.




Rock magnetism A difference in the magnetic mineralogy of certain flows could indicate a chemical remagnetization representing the local geomagnetic field at a later date leading to the more southerly VGP. However, the specimens in the two VGP populations encompass the same range of demagnetization behaviors. Therefore a rock magnetic explanation for the divergent directions is not readily apparent within the data. The demagnetization data (averaged by flow) of both high and low latitude VGPs are presented below.

```
In [16]: high_lat_sites = SLB_Data_hi_lat.er_site_names.tolist()
         low_lat_sites = SLB_Data_low_lat.er_site_names.tolist()


In [17]: ipmag.demag_magic('../Data/SLB/', plot_by='site', treat='T',
           individual=low_lat_sites, average_measurements=True,
           single_plot=True)

8204 records read from ../Data/SLB/magic_measurements.txt
```



```
In [18]: ipmag.demag_magic('../Data/SLB/', plot_by='site', treat='T',
                           individual=high_lat_sites, average_measurements=True,
                           single_plot=True)
```



8204 records read from ../Data/SLB/magic_measurements.txt



Patterns in the distribution and/or secular variation of poles Considered individually, both VGP populations are Fisher-distributed.


In [19]: `ipmag.fishqq(SLB_Data_low_lat['vgp_lon'].tolist(), SLB_Data_low_lat['vgp_lat'].tolist())`


Out[19]: {
 'Dec': 189.87618031371633,
 'Inc': 9.225749891843499,
 'Me': 0.77027561013556023,
 'Me_critical': 1.094,
 'Mode': 'Mode 1',
 'Mu': 0.93893808899861431,
 'Mu_critical': 1.207,
 'N': 10,
 'Test_result': 'consistent with Fisherian model'}



In [20]: ipmag.fishqq(SLB_Data_hi_lat['vgp_lon'].tolist(), SLB_Data_hi_lat['vgp_lat'].tolist())

```
Out[20]: {'Dec': 189.43399983176758,
'Inc': 32.306644938683029,
'Me': 1.1071268307909761,
'Me_critical': 1.094,
'Mode': 'Mode 1',
'Mu': 0.60645097771482315,
'Mu_critical': 1.207,
'N': 30,
'Test_result': 'Fisherian model rejected'}
```






One last way to approach this problem is to inspect the path of the paleomagnetic poles being traced out by secular variation. We can look for potential patterns in the path of the poles through time by considering each cooling unit's stratigraphic placement.

```
In [21]: # upload strat heights
SLB_strat_df = pd.read_csv('../Data/SLB/er_sites.txt', sep='\t', skiprows=1,
                           usecols=['er_site_name', 'site_height'])
SLB_strat_df = SLB_strat_df.set_index('er_site_name')

m = pole_figure_appearance()

centerlon, centerlat = m(SLB_Data['vgp_lon'].tolist(), SLB_Data['vgp_lat'].tolist())
m.plot(centerlon, centerlat, alpha=0.5, zorder=1)
m.scatter(centerlon, centerlat, marker='.', s=300.0,
          c = SLB_strat_df.site_height.tolist(), cmap='Paired', zorder=2)

plt.colorbar(label='Stratigraphy (meters)', shrink=0.7)
plt.show()
```



From this view, we notice the relatively tight grouping of poles among certain stratigraphic intervals as they trace out sections of the entire distribution. However, there is still no apparent explanation for the occasional excursion of the geomagnetic pole to lower latitudes. It therefore seems likely that the divergence of these two VGP populations can be attributed to a recurrent geomagnetic excursion rather than a post-emplacement offset of magnetizations. We also note the similarity of this anomalous behavior to the distribution of paleomagnetic poles from the Lake Shore Traps, as discussed by Diehl and Haig (1994) (see *Lake Shore Traps* section below).

2.3 Combine new SLB data with Tauxe and Kodama (2009)


```
In [22]: combined_SLB_lon = NSVG_ns1['pole_lon'].tolist() + SLB_Data['vgp_lon'].tolist()
combined_SLB_lat = NSVG_ns1['pole_lat'].tolist() + SLB_Data['vgp_lat'].tolist()
combined_SLB_mean = ipmag.fisher_mean(combined_SLB_lon, combined_SLB_lat)
```

```
In [23]: m = pole_figure_appearance()
ipmag.plot_pole(m, NSVG_ns1_mean['dec'], NSVG_ns1_mean['inc'],
                 NSVG_ns1_mean['alpha95'], marker='s', color='b',
                 label='NSVG mean (nswu; Tauxe and Kodama (2009))')
ipmag.plot_vgp(m, combined_SLB_lon, combined_SLB_lat, color='g',
                 label='combined SLB (this study and Tauxe and Kodama (2009))')
ipmag.plot_vgp(m, NSVG_ns1['pole_lon'].tolist(), NSVG_ns1['pole_lat'].tolist(),
                 color='springgreen', label='SLB (ns1; Tauxe and Kodama (2009))')
ipmag.plot_pole(m, combined_SLB_mean['dec'], combined_SLB_mean['inc'],
                 combined_SLB_mean['alpha95'], marker='s', color='g')
```

```
ipmag.print_pole_mean(combined_SLB_mean)

plt.legend()
plt.savefig('~/Code_output/NSVG_SLB_poles_combined.pdf')
plt.show()
```

Plong: 187.8 Plat: 27.1
Number of directions in mean (n): 50
Angular radius of 95% confidence (A_95): 3.0
Precision parameter (k) estimate: 46.5





This combined mean still reflects a non-Fisherian distribution, as seen below.

In [24]: ipmag.fishqq(combined_SLB_lon,combined_SLB_lat)

Out[24]: {'Dec': 187.79191234206988,
'Inc': 27.199685356762615,

```
'Me': 0.53556909069644332,  
'Me_critical': 1.094,  
'Mode': 'Mode 1',  
'Mu': 1.4805683534761531,  
'Mu_critical': 1.207,  
'N': 50,  
'Test_result': 'Fisherian model rejected'}
```





3 Lake Shore Traps

We first import paleomagnetic from Diehl and Haig (1994).

```
In [25]: Diehl1994a_LST_Data_all=pd.read_csv('../Data/Previous_studies/Diehl1994a_data.csv',sep=',')
#Kulakov2013 reported data for the flow LST28 that supersedes
#the Diehl direction which should accordingly be dropped
Diehl1994a_LST_Data=Diehl1994a_LST_Data_all.drop(17)
Diehl1994a_LST_Data.reset_index(inplace=True)
Diehl1994a_LST_Data.head()
```

This file can be found in the Data Repository under /Data/Previous_studies/Diehl1994a_data.csv.

Next, we import paleomagnetic data from Kulakov et al. (2013). The original Kulakov et al. (2013) data set incorporated the data of Diehl and Haig (1994); however, we elect to manually combine these data and recalculate the associated grand mean here.


```
In [26]: Kulakov2013a_LST_Data=pd.read_csv('../Data/Previous_studies/Kulakov2013a_data.csv',sep=',')
Kulakov2013a_LST_Data.head()
```

This file can be found in the Data Repository under /Data/Previous_studies/Kulakov2013a_data.csv.

```
In [27]: LST_Diehl_VGPs=ipmag.make_di_block(Diehl1994a_LST_Data['vgp_lon'],
                                            Diehl1994a_LST_Data['vgp_lat'])
```

```
LST_Kulakov_VGPs=ipmag.make_di_block(Kulakov2013a_LST_Data['vgp_lon'],
```



```
Kulakov2013a_LST_Data['vgp_lat'])  
  
LST_VGPs=np.concatenate((LST_Diehl_VGPs,LST_Kulakov_VGPs))  
  
LST_Diehl_mean = pmag.fisher_mean(LST_Diehl_VGPs)  
LST_Kulakov_mean=pmag.fisher_mean(LST_Kulakov_VGPs)  
LST_all_mean=pmag.fisher_mean(LST_VGPs)  
  
In [28]: m = Basemap(projection='ortho',lat_0=35,lon_0=200,resolution='c',  
                     area_thresh=50000)  
pole_figure_appearance()  
  
ipmag.plot_vgp(m,Diehl1994a_LST_Data['vgp_lon'].tolist(),  
                Diehl1994a_LST_Data['vgp_lat'].tolist(),  
                label='LST dikes VGPs (Diehl1994a)',color='r')  
ipmag.plot_vgp(m,Kulakov2013a_LST_Data['vgp_lon'].tolist(),  
                Kulakov2013a_LST_Data['vgp_lat'].tolist(),  
                label='LST dikes VGPs (Kulakov2013a)',color='b')  
ipmag.plot_pole(m,LST_all_mean['dec'],LST_all_mean['inc'],  
                 LST_all_mean['alpha95'], label='LST dikes mean pole',marker='s')  
ipmag.plot_pole(m,LST_Diehl_mean['dec'],LST_Diehl_mean['inc'],  
                 LST_Diehl_mean['alpha95'], color='r',  
                 label='Diehl1994a mean',marker='s')  
#from Kulakov2013a  
ipmag.plot_pole(m,186.4,23.1,4.0,color='g',  
                 label='mean as calculated in Kulakov2013a',marker='s')  
  
plt.legend(loc=4)  
plt.savefig('Code_output/LST_vgps.pdf')  
plt.show()
```



Here we test whether the distribution of Lake Shore Traps VGPs are Fisherian.


In [29]: `ipmag.fishqq(di_block=LST_VGPs)`

Out[29]: {
 'Dec': 185.61848690207148,
 'Inc': 23.207239038463833,
 'Me': 1.1080580586218494,
 'Me_critical': 1.094,
 'Mode': 'Mode 1',
 'Mu': 1.8795789739707827,
 'Mu_critical': 1.207,
 'N': 49,
 'Test_result': 'Fisherian model rejected'}



Here we test for a common mean between the Lake Shore Traps VGPs and our newly combined data from the Schroeder-Lutsen basalts. As seen below, although the A_{95} ellipses overlap, the two means can (barely) be distinguished at the 95% confidence level by the criteria of the Watson V test, which calculates Watson's V statistic ([Watson, 1956](#)) from input files through Monte Carlo simulation and to tests whether two populations of directional data could have been drawn from a common mean. The critical angle between the two sample mean directions and the corresponding [McFadden and McElhinny \(1990\)](#) classification is printed below.

```
In [30]: ipmag.common_mean_bootstrap(LST_VGPs,ipmag.make_di_block(combined_SLB_lon,combined_SLB_lat))
```



```
In [31]: ipmag.common_mean_watson(LST_VGPs,
                                 ipmag.make_di_block(combined_SLB_lon,combined_SLB_lat),
                                 plot='yes')
```

Results of Watson V test:

Watson's V: 6.0


Critical value of V: 5.9

"Fail": Since V is greater than V_{crit}, the two means can be distinguished at the 95% confidence level.

M&M1990 classification:

Angle between data set means: 4.4

Critical angle for M&M1990: 4.4



4 Michipicoten Island

4.1 Palmer and Davis (1987) data

```
In [32]: Palmer1987_data=pd.read_csv('../Data/Previous_studies/Palmer1987a_data_combined_sites.csv')
         ipmag.vgp_calc(Palmer1987_data)
```

```
Out[32]: View data of Palmer and Davis (1987) in the MagIC database
          (http://earthref.org/magic/m006169dt20061219133107)
```

The code below splits this dataframe into sites from older main stage basalts that have been correlated to the Mamainse Point Formation, sites from the intrusive suite, and sites from the Michipicoten Island Formation.

While not explicitly described as such, it appears that each of the sites sampled in Palmer and Davis (1987) are treated as single cooling units except for site 'KM' which is described as being collected over a stratigraphic interval containing multiple flows. As the site is not from a single cooling unit, data from this are excluded. Also, site 18 is described as coming from a fault bound block with anomalous steep dip where a structural attitude could not be determined and should be excluded.

Palmer and Davis (1987) sampled the Channel Lake Member, the Cuesta Member, and the Davieaux Island Member at multiple sites from which VGPs were found and used in the final calculation of the Michipicoten Island Formation mean paleomagnetic pole. However, field observations reveal that some of these sites represented the redundant sampling of a single cooling unit, meaning that several VGPs that were originally treated as independent should have been combined with one another in the final calculation. Here we reduce the Palmer and Davis (1987) data set to satisfy these concerns. The reduced number of sites for each Michipicoten Island Formation member is shown below.

```
In [33]: print 'Number of Cuesta Member flows: ',
        len(Palmer1987_data.loc[Palmer1987_data['member']=='Cuesta_Member'])
print 'Number of Channel Lake Member flows: ',
        len(Palmer1987_data.loc[Palmer1987_data['member']=='Channel_Lake_Member'])
print 'Number of Quebec Harbour Member flows: ',
        len(Palmer1987_data.loc[Palmer1987_data['member']=='Quebec_Harbour_Member'])
print 'Number of South Shore Member flows: ',
        len(Palmer1987_data.loc[Palmer1987_data['member']=='South_Shore_Member'])
print 'Number of Davieaux Island Member flows: ',
```

```

len(Palmer1987_data.loc[Palmer1987_data['member']=='Davieaux_Island_Member'])
print 'Total number of cooling units from the Michipicoten Island Formation: ',
len(Palmer1987_data.loc[Palmer1987_data['formation']=='Michipicoten_Island'])

Number of Cuesta Member flows:  2
Number of Channel Lake Member flows:  1
Number of Quebec Harbour Member flows:  1
Number of South Shore Member flows:  3
Number of Davieaux Island Member flows:  1
Total number of cooling units from the Michipicoten Island Formation:  8

In [34]: MI_Mamainse_Fm=Palmer1987_data.ix[Palmer1987_data['formation'] == 'Quebec_Mine']
MI_Mamainse_Fm.reset_index(inplace=True, drop=True)
#make dataframe that excludes sites 18 and KM and reset the index of the dataframe
#MI_Mamainse_Fm=MI_Mamainse_Fm_all.drop([1,8])
MI_Mamainse_Fm.reset_index(inplace=True)
MI_Mamainse_Fm_VGPs = ipmag.make_di_block(MI_Mamainse_Fm['vgp_lon'],
                                             MI_Mamainse_Fm['vgp_lat'])
MI_Mamainse_Fm_mean = pmag.fisher_mean(MI_Mamainse_Fm_VGPs)
print 'Michipicoten Island Mamainse Formation mean:'
ipmag.print_pole_mean(MI_Mamainse_Fm_mean)

MI_intrusions=Palmer1987_data.ix[Palmer1987_data['formation'] == 'intrusive']
MI_intrusions.reset_index(inplace=True, drop=True)
MI_intrusions_VGPs = ipmag.make_di_block(MI_intrusions['vgp_lon'],
                                           MI_intrusions['vgp_lat'])
MI_intrusions_mean = pmag.fisher_mean(MI_intrusions_VGPs)
print '\nMichipicoten Island intrusions mean:'
ipmag.print_pole_mean(MI_intrusions_mean)

Michipicoten_Island_Fm=Palmer1987_data.ix[Palmer1987_data['formation'] ==
                                           'Michipicoten_Island']
Michipicoten_Island_Fm.reset_index(inplace=True, drop=True)
Michipicoten_Island_Fm_VGPs = ipmag.make_di_block(Michipicoten_Island_Fm['vgp_lon'],
                                                 Michipicoten_Island_Fm['vgp_lat'])
Michipicoten_Island_Fm_mean = pmag.fisher_mean(Michipicoten_Island_Fm_VGPs)
print '\nMichipicoten Island Formation mean:'
ipmag.print_pole_mean(Michipicoten_Island_Fm_mean)

m = pole_figure_appearance()

ipmag.plot_vgp(m,MI_Mamainse_Fm['vgp_lon'].tolist(),MI_Mamainse_Fm['vgp_lat'].tolist(),
               label='MI_MP VGPs',color='g')
ipmag.plot_pole(m,MI_Mamainse_Fm_mean['dec'],MI_Mamainse_Fm_mean['inc'],
                MI_Mamainse_Fm_mean['alpha95'],
                label='MI_MP_mean',marker='s',color='g')
ipmag.plot_pole(m,183.2,31.2,2.5,label='upper Mamainse (S-H et al., 2014)',marker='s')

ipmag.plot_vgp(m,MI_intrusions['vgp_lon'].tolist(),MI_intrusions['vgp_lat'].tolist(),
               label='intrusion VGPs',color='blue')
ipmag.plot_pole(m,MI_intrusions_mean['dec'],MI_intrusions_mean['inc'],
                MI_intrusions_mean['alpha95'],
                label='intrusion mean',marker='s',color='blue')

ipmag.plot_vgp(m,Michipicoten_Island_Fm['vgp_lon'].tolist(),

```

```
Michipicoten_Island_Fm['vgp_lat'].tolist(),
label='Mich. Fm VGPs',color='goldenrod')
ipmag.plot_pole(m,Michipicoten_Island_Fm_mean['dec'],Michipicoten_Island_Fm_mean['inc'],
Michipicoten_Island_Fm_mean['alpha95'],
label='Mich. Fm mean',marker='s',color='goldenrod')
plt.legend()
plt.show()
```

Michipicoten Island Mamainse Formation mean:

Plong: 185.6 Plat: 36.9

Number of directions in mean (n): 7

Angular radius of 95% confidence (A_95): 13.4

Precision parameter (k) estimate: 21.2

Michipicoten Island intrusions mean:

Plong: 165.7 Plat: 23.9

Number of directions in mean (n): 3

Angular radius of 95% confidence (A_95): 22.5

Precision parameter (k) estimate: 31.2


Michipicoten Island Formation mean:

Plong: 174.9 Plat: 25.5

Number of directions in mean (n): 8

Angular radius of 95% confidence (A_95): 7.7

Precision parameter (k) estimate: 53.2



In [35]: Michipicoten_Island_Fm

4.2 New data from the Michipicoten Island Formation

New paleomagnetic data from the Michipicoten Island Formation are imported below. We first separate data by *in situ* (“geo”) and tilt-corrected (“tc”) coordinates. We then further separate data by member (“SS” = South Shore Member basalts, “CM” = Cuesta Member andesite).

```
In [36]: All_Michi = pd.read_csv('../Data/Michipicoten/pmag_results.txt',
                               sep='\t', skiprows=1)
All_Michi_geo = All_Michi.ix[All_Michi['pole_comp_name'] == 'HT']\n
               .ix[All_Michi['tilt_correction']==0]
All_Michi_tc = All_Michi.ix[All_Michi['pole_comp_name'] == 'HT']\n
               .ix[All_Michi['tilt_correction']==100]
All_Michi_tc.reset_index(inplace=True, drop=True)
All_Michi_geo.reset_index(inplace=True, drop=True)
```

Out[36]: View raw data in the MagIC contribution of this study
(<https://earthref.org/MagIC/doi/10.1130/L580.1>)

If you are having trouble opening the file via this link, the file can be found in the Data Repository under /Data/Michipicoten/pmag_results.

```
In [37]: SS_final = All_Michi.ix[All_Michi['er_site_names'].str.startswith('SS')]
SS_final_tc = All_Michi_tc.ix[All_Michi_tc['er_site_names'].str.startswith('SS')]
SS_final_geo = All_Michi_geo.ix[All_Michi_geo['er_site_names'].str.startswith('SS')]
SS_final_tc.head()
```

```
In [38]: CM_final_tc = All_Michi_tc.ix[All_Michi_tc['er_site_names'].str.startswith('CM')]
CM_final_tc
```

Paleomagnetic site means of the South Shore basalts are plotted below with the overall mean.

```
In [39]: plt.figure(num=1,figsize=(7,7))
ipmag.plot_net(1)

SS_dec = SS_final_tc['average_dec'].tolist()
SS_inc = SS_final_tc['average_inc'].tolist()
SS_a95 = SS_final_tc['average_alpha95'].tolist()


for i in range(len(SS_dec)):
    ipmag.plot_di_mean(SS_dec[i], SS_inc[i], SS_a95[i])

SS_mean_dir = ipmag.fisher_mean(SS_dec,SS_inc)

ipmag.plot_di_mean(SS_mean_dir['dec'],SS_mean_dir['inc'],
                    SS_mean_dir['alpha95'],marker='s', color='r')
ipmag.print_direction_mean(SS_mean_dir)
plt.savefig('Code_output/All_SS_data.pdf')
plt.title('Magnetization directions of South Shore basalts')
plt.clabel
plt.show()
```

```
Dec: 287.6  Inc: 10.4
Number of directions in mean (n): 21
Angular radius of 95% confidence (a_95): 5.4
Precision parameter (k) estimate: 35.4
```



Magnetization directions of South Shore basalts



A Fisher Q-Q test reveals these data to be consistent with the Fisherian model.

In [40]: `ipmag.fishqq(SS_dec, SS_inc)`

Out [40]: {
 'Dec': 287.60360855657012,
 'Inc': 10.297033320865525,
 'Me': 0.92123018046225358,
 'Me_critical': 1.094,
 'Mode': 'Mode 1',
 'Mu': 0.92607542329666181,
 'Mu_critical': 1.207,
 'N': 21,
 'Test_result': 'consistent with Fisherian model'}



The bedding is similar through the South Shore Member such that a fold test is inconclusive (see below). A more complete treatment of our structural analysis of the Michipicoten Island lava flows, including a walkthrough of how we determined tilt-correction for different stratigraphic sections of the South Shore Member, is included within the Data Repository as a separate Jupyter notebook.

```
In [41]: # get bedding info
bedding = pd.read_csv('../Data/Michipicoten/er_samples.txt',
                      sep='\t', skiprows=1, usecols=['er_site_name',
                      'sample_bed_dip',
                      'sample_bed_dip_direction'])

In [42]: bedding = bedding.drop_duplicates()
bedding = bedding.set_index('er_site_name')

In [43]: bedding.head()

Out[43]:
er_site_name  sample_bed_dip  sample_bed_dip_direction
CM1           31.8            171.9
CM2           25.5            177.7
SS1           20.0            237.6
SS10          20.0            237.6
SS11          20.0            237.6


In [44]: SS_diddd = []

for site in bedding.index:
    if str(site) in SS_final_geo.er_site_names.tolist():
        SS_diddd.append([float(SS_final_geo.loc[SS_final_geo['er_site_names']\
                                              ==str(site)]['average_dec']),
                          float(SS_final_geo.loc[SS_final_geo['er_site_names']\
                                              ==str(site)]['average_inc']),
                          bedding.loc[str(site), 'sample_bed_dip_direction'],
                          bedding.loc[str(site), 'sample_bed_dip']])


In [45]: ipmag.bootstrap_fold_test(np.array(SS_diddd))

doing 1000 iterations...please be patient...
```

Geographic



Tilt-corrected




tightest grouping of vectors obtained at (95% confidence bounds):

7 - 95 percent unfolding

range of all bootstrap samples:

-10 - 119 percent unfolding




Paleomagnetic site means of the Cuesta Member andesite are plotted below. Only two Cuesta Member flows were identified in the field, so a Cuesta Member mean would not have much significance.

```
In [46]: plt.figure(num=1,figsize=(5,5))
ipmag.plot_net(1)

CM_dec = CM_final_tc['average_dec'].tolist()
CM_inc = CM_final_tc['average_inc'].tolist()
CM_a95 = CM_final_tc['average_alpha95'].tolist()

for i in range(len(CM_dec)):
    ipmag.plot_di_mean(CM_dec[i], CM_inc[i], CM_a95[i])

plt.show()
```



4.3 Combine new Michipicoten data with Palmer and Davis (1987)

Given that we have obtained U-Pb dates for the West Sand Bay tuff and the Davieaux Island Member rhyolite, we are interested in generating a pole that is bracketed by these dates. The units between and including these dates are the West Sand Bay Member, the Quebec Harbor Member, the East Sand Bay Member, the South Shore Member and the Davieaux Island Member. Therefore we combine our data from the South Shore Member with the Quebec Harbour Member and Davieaux Island Member data from Palmer and Davis (1987).

```
In [47]: Michipicoten_Island_Fm.iloc[[3,7], :]

In [48]: Palmer_data_trimmed = Michipicoten_Island_Fm.iloc[[3,7], :]
combined_Michi_lon = Palmer_data_trimmed['vgp_lon'].tolist() \
+ SS_final_tc['vgp_lon'].tolist()
combined_Michi_lat = Palmer_data_trimmed['vgp_lat'].tolist() \
+ SS_final_tc['vgp_lat'].tolist()
combined_Michi_mean = ipmag.fisher_mean(combined_Michi_lon, combined_Michi_lat)

In [49]: SS_final_mean = ipmag.fisher_mean(SS_final_tc['vgp_lon'].tolist(),
SS_final_tc['vgp_lat'].tolist())
```

```
m = pole_figure_appearance()

ipmag.plot_vgp(m,SS_final_tc['vgp_lon'].tolist(),
                SS_final_tc['vgp_lat'].tolist(),
                label='South Shore Member VGP (this study)',color='b')

ipmag.plot_vgp(m,Palmer_data_trimmed['vgp_lon'].tolist(),
                Palmer_data_trimmed['vgp_lat'].tolist(),
                label='Palmer and Davis (1987) VGP used in mean pole',
                color='purple')

ipmag.plot_pole(m,combined_Michi_mean['dec'],combined_Michi_mean['inc'],
                 combined_Michi_mean['alpha95'], label='Michipicoten Island Fm. \
grand mean (this study and Palmer and Davis, 1987)',
                 marker='s',markersize=25, color='red')

plt.legend()
plt.savefig('Code_output/Michi_VGPs.pdf')


print "South Shore Member pole: "
ipmag.print_pole_mean(SS_final_mean)

print "\nGrand mean: "
ipmag.print_pole_mean(combined_Michi_mean)

plt.show()



South Shore Member pole:
Plong: 174.7 Plat: 15.8
Number of directions in mean (n): 21
Angular radius of 95% confidence (A_95): 4.3
Precision parameter (k) estimate: 55.4

Grand mean:
Plong: 174.7 Plat: 17.0
Number of directions in mean (n): 23
Angular radius of 95% confidence (A_95): 4.4
Precision parameter (k) estimate: 48.4
```



In [50]: `ipmag.fishqq(combined_Michi_lon,combined_Michi_lat)`

Out[50]: {
 'Dec': 174.68056363405637,
 'Inc': 16.974664865055974,
 'Me': 0.8235241344357922,
 'Me_critical': 1.094,
 'Mode': 'Mode 1',
 'Mu': 1.1041506813063025,
 'Mu_critical': 1.207,
 'N': 23,
 'Test_result': 'consistent with Fisherian model'}



5 Paleogeography of Laurentia

5.1 Late Rift Poles Compilation

Here we compile selected paleomagnetic poles of the Keweenawan Track and add our newly developed data. This section of the code integrates the functionality of the GPlates software for our paleogeographic reconstruction of Laurentia. The `pygplates` package is a dependency of the following code blocks and is imported below. If you wish to render this notebook interactively, the `pygplates` package can be downloaded from the GPlates website: <http://www.gplates.org/download.html>

```
In [51]: # If this code block gives an error after you have already
# installed PyGPlates, you probably still need to add the package to your path
import pygplates

In [52]: # 1080 Ma = Nonesuch pole in recon, 1070 Ma = Freda pole, 1050 = Jacobsville
for reconstruction_time in [1109, 1105, 1100, 1097, 1090, 1083, 1080, 1070, 1050]:
    rotation_model = pygplates.RotationModel(
        '../Data/Reconstruction/Late_Rift_recon.rot')
    plate = pygplates.FeatureCollection(
        '../Data/Reconstruction/1000_Laurentia.gpml')
    plate_outline = pygplates.FeatureCollection(
        '../Data/Reconstruction/1000_Laurentia2.gpml')
    export_filename = './Code_output/Reconstructions/reconstructed_{0}Ma.shp'
        .format(reconstruction_time)
    export_outline_filename =
        './Code_output/Reconstructions/reconstructed_outline_{0}Ma.shp'
        .format(reconstruction_time)
    pygplates.reconstruct(plate, rotation_model, export_filename,
                          reconstruction_time)
    pygplates.reconstruct(plate_outline, rotation_model,
                          export_outline_filename, reconstruction_time)

In [53]: fig = plt.figure(figsize=(12,6))
ax1 = fig.add_subplot(121)
m = Basemap(projection='ortho',lat_0=35,lon_0=200,resolution='c',
            area_thresh=50000)
ax1.set_aspect(30, adjustable='box')
m.drawcoastlines(linewidth=0.25)
m.fillcontinents(color='bisque',lake_color='white')
m.drawmapboundary(fill_color='white')
m.drawmeridians(np.arange(0,360,30))
m.drawparallels(np.arange(-90,90,30))
m.readshapefile('../Data/Reconstruction/Laurentia', 'Laurentia', linewidth=1)
# Swanson-Hysell et al., 2014a -- 1100 Ma
ipmag.plot_pole(m, 227.0, 49.5, 5.3, marker='s',
                 label='Mamainse Pt. (lowermostR)')
# Swanson-Hysell et al., 2014b -- 1105 Ma
ipmag.plot_pole(m, -158.3, 42.5, 3.7, marker='s',
                 label='Osler Volcanic Group (upperR)')
# Swanson-Hysell et al., 2014a -- 1100 Ma
ipmag.plot_pole(m, -170.3, 36.1, 4.9, marker='s',
                 label='Mamainse Pt. (lowerN, upperR)')
# Tauxe and Kodama (2009) -- 1097 Ma
ipmag.plot_pole(m,NSVG_nswu_mean['dec'],NSVG_nswu_mean['inc'],
                 NSVG_nswu_mean['alpha95'], marker='s',label='NSVG')
```

```

# this study + Tauxe and Kodama (2009) -- 1090 Ma
ipmag.plot_pole(m,combined_SLB_mean['dec'],combined_SLB_mean['inc'],
                 combined_SLB_mean['alpha95'], marker='s',
                 label='SLB')
# Combine Books (1972) and Hnat et al. (2006) -- 1093 Ma
ipmag.plot_pole(m, 182.0, 27.1, 2.2, marker='s',
                 label='Portage Lake Volcanics')
# Kulakov et al. (2013)
ipmag.plot_pole(m,LST_all_mean['dec'],LST_all_mean['inc'],
                 LST_all_mean['alpha95'], label='LST dikes mean pole',
                 marker='s')
# this study + Palmer and Davis (1987) -- 1083 Ma
ipmag.plot_pole(m,combined_Michi_mean['dec'],combined_Michi_mean['inc'],
                 combined_Michi_mean['alpha95'],
                 label='Michipicoten Island Fm.',marker='s')
# Henry et al. (1977)
ipmag.plot_pole(m, 178.1, 7.6, 5.5, label='Nonesuch Shale',
                 marker='s')
# Henry et al. (1977)
ipmag.plot_pole(m, 179, 2.2, 4.2,label='Freda Sandstone',
                 marker='s')
# Roy et al. (1978)
ipmag.plot_pole(m, 184, -10, 4.2,label='Jacobsville Sandstone',
                 marker='s')


ax1.legend(fontsize=8, loc='lower right')

ax2 = fig.add_subplot(122)
m = Basemap(projection='ortho', lat_0=30, lon_0=-16)
m.drawmapboundary(fill_color='white')
m.drawmeridians(np.arange(0,360,30))
m.drawparallels(np.arange(-90,90,30))
m.readshapefile('./Code_output/Reconstructions/reconstructed_outline_{0}Ma'
                .format(1109),
                'Mamainse_lowermost', color='navy', linewidth=1)
patches = []
for info, shape in zip(m.Mamainse_lowermost_info, m.Mamainse_lowermost):
    patches.append(Polygon(np.array(shape), True))
ax2.add_collection(PatchCollection(patches, facecolor= 'navy', edgecolor='k',
                                    linewidths=1., zorder=2))
m.readshapefile('./Code_output/Reconstructions/reconstructed_{0}Ma'.format(1109),
                'Mamainse_lowermost', color='k', linewidth=1)
m.readshapefile('./Code_output/Reconstructions/reconstructed_outline_{0}Ma'
                .format(1105),
                'Osler_outline', color='slategrey', linewidth=1)
patches = []
for info, shape in zip(m.Osler_outline_info, m.Osler_outline):
    patches.append(Polygon(np.array(shape), True))
ax2.add_collection(PatchCollection(patches, facecolor= 'slategrey',
                                    edgecolor='k', linewidths=1., zorder=2))
m.readshapefile('./Code_output/Reconstructions/reconstructed_{0}Ma'.format(1105),
                'Osler', color='k', linewidth=1)
m.readshapefile('./Code_output/Reconstructions/reconstructed_outline_{0}Ma'
                .format(1100),

```

```
        'Mamainse', color='lightgray', linewidth=1)
patches = []
for info, shape in zip(m.Mamainse_info, m.Mamainse):
    patches.append( Polygon(np.array(shape), True) )
ax2.add_collection(PatchCollection(patches, facecolor= 'lightgray',
                                    edgecolor='k', linewidths=1., zorder=2))
m.readshapefile('./Code_output/Reconstructions/reconstructed_{0}Ma'
                .format(1100),
                'Mamainse', color='k', linewidth=1)
m.readshapefile('./Code_output/Reconstructions/reconstructed_outline_{0}Ma'
                .format(1097),
                'NSVG', color='b', linewidth=1)
patches = []
for info, shape in zip(m.NSVG_info, m.NSVG):
    patches.append( Polygon(np.array(shape), True) )
ax2.add_collection(PatchCollection(patches, facecolor= 'b', edgecolor='k',
                                    linewidths=1., zorder=2))
m.readshapefile('./Code_output/Reconstructions/reconstructed_{0}Ma'.format(1097),
                'NSVG', color='k', linewidth=1)
m.readshapefile('./Code_output/Reconstructions/reconstructed_outline_{0}Ma'
                .format(1090),
                'SLB', color='g', linewidth=1)
patches = []
for info, shape in zip(m.SLB_info, m.SLB):
    patches.append( Polygon(np.array(shape), True) )
ax2.add_collection(PatchCollection(patches, facecolor= 'g', edgecolor='k',
                                    linewidths=1., zorder=2))
m.readshapefile('./Code_output/Reconstructions/reconstructed_{0}Ma'.format(1090),
                'SLB', color='k', linewidth=1)
m.readshapefile('./Code_output/Reconstructions/reconstructed_outline_{0}Ma'
                .format(1083),
                'Michipicoten', color='r', linewidth=1)
patches = []
for info, shape in zip(m.Michipicoten_info, m.Michipicoten):
    patches.append( Polygon(np.array(shape), True) )
ax2.add_collection(PatchCollection(patches, facecolor= 'r', edgecolor='k',
                                    linewidths=1., zorder=2))
m.readshapefile('./Code_output/Reconstructions/reconstructed_{0}Ma'.format(1083),
                'Michipicoten', color='k', linewidth=1)
m.readshapefile('./Code_output/Reconstructions/reconstructed_outline_{0}Ma'
                .format(1080),
                'Nonesuch', color='magenta', linewidth=1)
patches = []
for info, shape in zip(m.Nonesuch_info, m.Nonesuch):
    patches.append( Polygon(np.array(shape), True) )
ax2.add_collection(PatchCollection(patches, facecolor= 'magenta',
                                    edgecolor='k', linewidths=1., zorder=2))
m.readshapefile('./Code_output/Reconstructions/reconstructed_{0}Ma'.format(1080),
                'Nonesuch', color='k', linewidth=1)
m.readshapefile('./Code_output/Reconstructions/reconstructed_outline_{0}Ma'
                .format(1070),
                'Freda', color='goldenrod', linewidth=1)
patches = []
for info, shape in zip(m.Freda_info, m.Freda):
```



```
patches.append( Polygon(np.array(shape), True) )
ax2.add_collection(PatchCollection(patches, facecolor= 'goldenrod',
                                    edgecolor='k', linewidths=1., zorder=2))
m.readshapefile('./Code_output/Reconstructions/reconstructed_{0}Ma'.format(1070),
                'Freda', color='k', linewidth=1)
# plt.savefig('Code_output/poles.pdf')
plt.show()
```



5.2 Rates of plate motion

5.2.1 Latitudinal motion implied by Osler and Michipicoten poles

- The paleolatitude for reference location (Duluth, MN) resulting from pole 1 (Osler Volcanic Group) is: 44.055°
- The paleolatitude for reference location (Duluth, MN) resulting from pole 2 (Michipicoten Island Formation) is: 10.194°
- The rate of paleolatitudinal change implied by the poles pairs is: 17.257 cm/yr





2.5^{th} percentile is: 14.518 cm/yr

50^{th} percentile is: 17.250 cm/yr

97.5^{th} percentile is: 20.359 cm/yr

5.2.2 Latitudinal motion implied by Mamainse lowerN/upperR and Michipicoten poles

- The paleolatitude for reference location (Duluth, MN) resulting from pole 1 (Mamainse Point lowerN/upperR) is: 32.862°
- The paleolatitude for reference location (Duluth, MN) resulting from pole 2 (Michipicoten Island Formation) is: 10.194°
- The rate of paleolatitudinal change implied by the poles pairs is: 14.942 cm/yr





2.5^{th} percentile is: 11.576 cm/yr

50^{th} percentile is: 14.932 cm/yr

97.5^{th} percentile is: 18.313 cm/yr

5.2.3 Latitudinal motion implied by North Shore Volcanic Group and Michipicoten poles

- The paleolatitude for reference location (Duluth, MN) resulting from pole 1 (North Shore Volcanic Group) is: 27.825°
- The paleolatitude for reference location (Duluth, MN) resulting from pole 2 (Michipicoten Island Formation) is: 10.194°
- The rate of paleolatitudinal change implied by the poles pairs is: 18.325 cm/yr



2.5^{th} percentile is: 13.929 cm/yr

50^{th} percentile is: 18.329 cm/yr

97.5^{th} percentile is: 22.773 cm/yr

5.3 Age of Keweenawan sediments

5.3.1 *Nonesuch Shale*

There is much evidence to support the deposition of the Nonesuch shale shortly after or perhaps during late stage volcanism in the Midcontinent Rift (see main text). However, the magnetization of this formation cannot be dated as definitively. Symons et al. (2013) suggested that the magnetization of the Nonesuch shale is secondary, likely a product of oxidation and mineralization in the Nonesuch Formation that postdates its deposition and subsequent burial by the Freda sandstone. They dated this magnetization at 1063 ± 8 Ma by projecting Laurentia's latitudinal rate of motion to the paleolatitude implied by the Nonesuch paleomagnetic pole. Their analysis assumes a rate of motion consistent with the previously hypothesized slowdown of Laurentia during the late stage of rifting (Davis and Green, 1997). However, our results suggest no significant change in Laurentia's motion over this period. Redoing the analysis of Symons et al. (2013) with our revised rate estimates yields an age of Nonesuch shale magnetization of approximately 1078 Ma (see below).

Create a motion path of Laurentia that captures the paleolatitude of Duluth, MN at 5 myr intervals through active rifting (1108-1083 Ma).


```
In [54]: # Duluth, MN
SeedPoint = (46.78,-92.1)
MovingPlate = 1000
RelativePlate = 0
times = np.arange(1083,1108,5)

# Create a motion path feature
digitisation_time = 0
seed_points_at_digitisation_time = pygplates.MultiPointOnSphere([SeedPoint])
motion_path_feature = pygplates.Feature.create_motion_path(
    seed_points_at_digitisation_time,
    times,
    valid_time=(1110, 1000),
    relative_plate=RelativePlate,
    reconstruction_plate_id = MovingPlate)

# Create the shape of the motion path
reconstruction_time = 1083
reconstructed_motion_paths = []
pygplates.reconstruct(motion_path_feature, rotation_model,
    reconstructed_motion_paths, reconstruction_time,
    reconstruct_type=pygplates.ReconstructType.motion_path)

# get the reconstructed coordinates into numpy arrays
for reconstructed_motion_path in reconstructed_motion_paths:
    trail = reconstructed_motion_path.get_motion_path().to_lat_lon_array()

In [55]: plt.plot(times,np.flipud(trail[:,0]))
plt.title('Paleolatitude of Duluth, MN')
plt.xlabel('Time (Ma)')
plt.ylabel('Latitude')
plt.gca().grid()
plt.gca().invert_xaxis()
plt.show()
```




The latitudinal change plotted above appears fairly constant. We therefore make a least squares fit to this data below.

```
In [56]: m, c = np.linalg.lstsq(np.vstack([times, np.ones(len(times))]).T,
                               np.flipud(trail[:,0]))[0]
        print 'Rate of Laurentia\'s latitudinal motion = %0.4f degrees/myr' % (m)
        print 'or %0.4f cm/yr' % (m*11.132)
```

Rate of Laurentia's latitudinal motion = 1.3767 degrees/myr
or 15.3254 cm/yr

```
In [57]: plt.plot(times,np.flipud(trail[:,0]))
        times_new = np.arange(1073,1108,5)
        plt.plot(times_new, m*times_new + c, 'r')
        # Paleolatitude of Nonesuch is 5.6 degrees
        plt.plot([1070,1105], [5.6,5.6])
        NS_inferred_age = (5.6-c)/m
        plt.title('Paleolatitude of Duluth, MN')
        plt.xlabel('Time (Ma)')
        plt.ylabel('Latitude')
        plt.gca().grid()
        plt.gca().invert_xaxis()
        plt.show()
```

Inferred age of Nonesuch Shale if Laurentia continued moving at the same rate after Midcontinent Rift volcanism ended:
1078.4340 Ma



5.3.2 Jacobsville Sandstone

New detrital zircon dates from the Jacobsville sandstone, which unconformably overlies the Freda sandstone, reveal a much younger age for this formation than estimated for the Orono Group sediments ($<959 \pm 19$ Ma; Malone et al., 2016). The paleolatitude of Laurentia implied by the Neoproterozoic APWP was then matched with that implied by the Jacobsville paleomagnetic pole, yielding a set of possible ages for Jacobsville deposition: ~ 780 -755, ~ 700 -610 or ~ 570 -555 Ma (Malone et al., 2016). Motivations for redoing this analysis with paleomagnetic poles from both the APWP of Laurentia and the APWP of Baltica (hypothesized to have a longstanding connection with Laurentia's northeastern margin throughout the Neoproterozoic; Pisarevsky et al., 2003; Li et al., 2008; Evans, 2009) are outlined in the main text (*Discussion: Age of Keweenawan sediments*).

We first upload paleomagnetic poles of Laurentia as compiled by the Nordic Supercontinent Workshop, Haraldvagen, Norway in 2014 (see references in Figure 7 of the manuscript).

```
In [58]: Laurentia_poles = pd.read_csv('../Data/Reconstruction/Laurentia_Poles.csv',
                                     usecols=['Formation', 'Terrane', 'AgeUpper',
                                              'AgeLower', 'AgeNominal', 'A95',
                                              'SLat', 'SLon', 'PLat', 'PLon',
                                              'RefLat', 'RefLon'])
Laurentia_poles = Laurentia_poles.sort_values(by='AgeNominal',
                                              ascending=False)
Laurentia_poles.reset_index(inplace=True, drop=True)
Laurentia_poles
```

These paleomagnetic data (as compiled by the 2014 Nordic Supercontinent Workshop, Haraldvagen, Norway, and with additional data from this study) can be found in the Data Repository under /Data/Reconstruction/Laurentia_Poles.csv.


The paleolatitude of Duluth, MN (“RefLat”, “RefLon”) is then calculated for each of these paleomagnetic poles and plotted below.

```
In [59]: Laurentia_poles['Latitude'] = pd.Series(90 - np.rad2deg(np.arccos\
    (np.sin(np.deg2rad(Laurentia_poles['RefLat'])))\*\
    *np.sin(np.deg2rad(Laurentia_poles['PLat'])))\*\
    +np.cos(np.deg2rad(Laurentia_poles['RefLat'])))\*\
    *np.cos(np.deg2rad(Laurentia_poles['PLat'])))\*\
    *np.cos(np.deg2rad(Laurentia_poles['RefLon'])-Laurentia_poles['PLon']))))

In [60]: # First we must tweak the format of the pole list
Laurentia_copy = pd.DataFrame(Laurentia_poles, columns=['PLat', 'PLon'])
Laurentia_copy = Laurentia_copy.rename_axis({'PLat': 'PLat_rot',
                                             'PLon': 'PLon_rot'},
                                             axis='columns')
Laurentia_poles['PLat_rot'] = Laurentia_copy.PLat_rot
Laurentia_poles['PLon_rot'] = Laurentia_copy.PLon_rot
Laurentia_poles['AgeNominal_neg'] = Laurentia_poles.AgeNominal.apply(np.negative)

# Plot the paleolatitude of Duluth using Laurentia poles
fig = plt.figure(figsize=(20,6))

ax1 = fig.add_subplot(131)
ax1.add_patch(mpatch.Rectangle((1300, -12.2), -1000, 8.4,
                               facecolor='r', alpha=0.2))
ax1.plot([-8.1]*2000, 'r-')
ax1.errorbar(Laurentia_poles['AgeNominal'].tolist(),
              Laurentia_poles['Latitude'].tolist(),
              yerr=Laurentia_poles['A95'].tolist(),
              xerr=(Laurentia_poles['AgeUpper']-Laurentia_poles['AgeLower'])/2)
ax1.text(1150, -6, 'Paleolat. from Jacobsville Sandstone',
         color='r', withdash=True)
# Cut off at Tonian/Cryogenian boundary
ax1.set_xlim(1200,717)
ax1.set_ylim(-90,90)
ax1.set_ylabel('Paleolatitude of Laurentia (Laurentia poles)')
ax1.set_xlabel('Age (Ma)')
plt.show()
```



Next, we upload paleomagnetic poles of Baltica as compiled by the Nordic Supercontinent Workshop, Haraldvangen, Norway in 2014 (see references in Figure 7 of the manuscript).

```
In [61]: Baltica_poles = pd.read_csv('../Data/Reconstruction/Baltica_Poles.csv',
                                    usecols=['Formation', 'Terrane', 'AgeUpper',
                                             'AgeLower', 'AgeNominal', 'A95',
                                             'SLat', 'SLon', 'PLat', 'PLon',
                                             'RefLat', 'RefLon'])
Baltica_poles = Baltica_poles.sort_values(by='AgeNominal', ascending=False)
Baltica_poles.reset_index(inplace=True, drop=True)
Baltica_poles
```

These paleomagnetic data (as compiled by the 2014 Nordic Supercontinent Workshop, Haraldvangen, Norway) can be found in the Data Repository under /Data/Reconstruction/Baltica_Poles.csv.

We then rotate these Baltica poles based on the reconstruction of Evans (2009) and do the same calculation for paleolatitude of Duluth, MN.

```
In [62]: Baltica_poles['PLat_rot'], Baltica_poles['PLon_rot'] = pmag.PTrot([81.5,
                                                               -110.0,
```

```

        -50.0],
        Baltica_poles.PLat.tolist(),
        Baltica_poles.PLon.tolist())
Baltica_poles['Latitude'] = pd.Series(90 - np.rad2deg(np.arccos\
(np.sin(np.deg2rad(Baltica_poles['RefLat'])))\*
*np.sin(np.deg2rad(Baltica_poles['PLat_rot'])))\*
+np.cos(np.deg2rad(Baltica_poles['RefLat'])))\*
*np.cos(np.deg2rad(Baltica_poles['PLat_rot'])))\*
*np.cos(np.deg2rad(Baltica_poles['RefLon'])-Baltica_poles['PLon_rot'])))))

```

We then combine the Laurentia and Baltica poles.

```

In [63]: combined_poles = Baltica_poles.append(Laurentia_poles)
combined_poles = combined_poles.sort_values(by='AgeNominal', ascending=False)
combined_poles = combined_poles.loc[combined_poles['AgeNominal']<=1200]
# Cut off at Tonian/Cryogenian boundary
combined_poles = combined_poles.loc[combined_poles['AgeNominal']>=717]
combined_poles.reset_index(inplace=True, drop=True)
combined_poles['AgeNominal_neg'] = combined_poles.AgeNominal.apply(np.negative)

In [64]: plt.figure(figsize=(13,8))
plt.subplot2grid((2,3), (0,1), colspan=2)
m = Basemap(projection='moll', lat_0=30, lon_0=210, resolution='c', area_thresh=50000)
m.readshapefile('../Data/Reconstruction/Laurentia_Baltica',
    'Laurentia_Baltica', drawbounds=True, linewidth=1)
m.drawcoastlines(linewidth=0.25)
m.fillcontinents(color='bisque', lake_color='white')
m.drawmapboundary(fill_color='white')
m.drawmeridians(np.arange(0,360,30))
m.drawparallels(np.arange(-90,90,30))
Laurentia_poles = Laurentia_poles.loc[Laurentia_poles['AgeNominal']<=1200]
Laurentia_poles = Laurentia_poles.loc[Laurentia_poles['AgeNominal']>=635]
Laurentia_poles.reset_index(inplace=True, drop=True)
centerlon, centerlat = m(Laurentia_poles['PLon_rot'].tolist(),
    Laurentia_poles['PLat_rot'].tolist())
for n in range(len(Laurentia_poles)):
    ipmag.plot_pole(m, Laurentia_poles['PLon_rot'][n],
                    Laurentia_poles['PLat_rot'][n],
                    Laurentia_poles['A95'][n])
ipmag.plot_pole(m, 184, -10, 4.2, marker='s', color='r', markersize=6.0)
m.scatter(centerlon, centerlat, c=Laurentia_poles['AgeNominal_neg'].tolist(),
          cmap='cubehelix', s=40, zorder=101)
plt.colorbar(orientation='horizontal', shrink=0.6)

ax1 = plt.subplot2grid((2,3), (0,0), colspan=1)
ax1.add_patch(mp.Rectangle((1300, -12.2), -1000, 8.4,
                           facecolor='r', alpha=0.2))
ax1.plot([-8.1]*2000, 'r-')
ax1.errorbar(Laurentia_poles['AgeNominal'].tolist(),
              Laurentia_poles['Latitude'].tolist(),
              yerr=Laurentia_poles['A95'].tolist(),
              xerr=(Laurentia_poles['AgeUpper']-Laurentia_poles['AgeLower'])/2, c='b')
ax1.text(1150, -6, 'Paleolat. from Jacobsville Sandstone',
         color='r', withdash=True)
ax1.text(900, 70, 'Laurentia poles')

```


```
ax1.set_xlim(1200,700)
ax1.set_ylim(-90,90)
ax1.set_ylabel('Paleolatitude of Duluth, MN')
ax1.set_xlabel('Age (Ma)')

m = Basemap(projection='moll',lat_0=30,lon_0=210,resolution='c',
             area_thresh=50000)
plt.subplot2grid((2,3), (1,1), colspan=2)
m.readshapefile('../Data/Reconstruction/Laurentia_Baltica',
                 'Laurentia_Baltica', drawbounds=True, linewidth=1)
m.drawcoastlines(linewidth=0.25)
m.fillcontinents(color='bisque',lake_color='white')
m.drawmapboundary(fill_color='white')
m.drawmeridians(np.arange(0,360,30))
m.drawparallels(np.arange(-90,90,30))
centerlon, centerlat = m(combined_poles['PLon_rot'].tolist(),
                           combined_poles['PLat_rot'].tolist())
for n in range(len(combined_poles)):
    ipmag.plot_pole(m, combined_poles['PLon_rot'][n],
                    combined_poles['PLat_rot'][n],
                    combined_poles['A95'][n])
ipmag.plot_pole(m, 184, -10, 4.2, marker='s', color='r', markersize=6.0)
m.scatter(centerlon, centerlat, c=combined_poles['AgeNominal_neg'].tolist(),
          cmap='cubebehelix', s=40, zorder=101)
plt.colorbar(orientation='horizontal', shrink=0.6)


ax2 = plt.subplot2grid((2,3), (1,0), colspan=1)
ax2.add_patch(mpolygon.Rectangle((1300, -12.2), -1000, 8.4,
                                 facecolor='r', alpha=0.2))
ax2.plot([-8.1]*2000, 'r-')
ax2.errorbar(combined_poles['AgeNominal'].tolist(),
              combined_poles['Latitude'].tolist(),
              yerr=combined_poles['A95'].tolist(),
              xerr=(combined_poles['AgeUpper']-
                     combined_poles['AgeLower'])/2, c='g')
for n in range(len(Laurentia_poles)):
    ax2.errorbar(Laurentia_poles['AgeNominal'][n],
                  Laurentia_poles['Latitude'][n],
                  yerr=Laurentia_poles['A95'][n],
                  xerr=(Laurentia_poles['AgeUpper'][n]-
                         Laurentia_poles['AgeLower'][n])/2, c='b')

ax2.text(1000, 70, 'Laurentia + Baltica poles')
ax2.set_xlim(1200,700)
ax2.set_ylim(-90,90)
ax2.set_ylabel('Paleolatitude of Duluth, MN')
ax2.set_xlabel('Age (Ma)')

plt.savefig('Code_output/Laur_Bal_poles.svg')
plt.show()
```



As seen below, the Jacobsville paleomagnetic pole aligns well with the Early Cambrian segment of Laurentia's APWP, suggesting a possible alternative Cambrian age. However, a Cambrian deposition of the Jacobsville seems unlikely when considering geological and fault relationships in the rift structure (see main text).



6 Map of late stage volcanics paleomagnetic sites

The following is an overview map of all paleomagnetic sites used in the preceding analysis.

```
In [65]: fig = plt.figure(figsize=(8,8))
m = Basemap(projection='merc',llcrnrlat=46.2,urcrnrlat=50,llcrnrlon=-92.5,
            urcrnrlon=-84,resolution='i',area_thresh = 0.1) #lat_ts=-25
m.drawrivers(color='#99ffff')
m.drawcoastlines()
#m.drawcountries(linewidth=1.5)
m.drawmapboundary(fill_color='#99ffff')
m.fillcontinents(color='#cc9966',lake_color='#99ffff')
parallels = np.arange(-90,90,2.)
m.drawparallels(parallels,labels=[1,0,0,0],fontsize=10)
meridians = np.arange(0.,360.,2.)
```


```
m.drawmeridians(meridians,labels=[0,0,0,1],fontsize=10)
plt.title('Late Stage paleomagnetic sites')

LST_site_lon=[]
LST_site_lat=[]
for n in range(0,len(Kulakov2013a_LST_Data)):
    LST_site_lon.append(Kulakov2013a_LST_Data['site_lon'][n])
    LST_site_lat.append(Kulakov2013a_LST_Data['site_lat'][n])
for n in range(0,len(Diehl1994a_LST_Data_all)):
    LST_site_lon.append(Diehl1994a_LST_Data_all['site_lon'][n])
    LST_site_lat.append(Diehl1994a_LST_Data_all['site_lat'][n])

LST_x,LST_y = m(LST_site_lon,LST_site_lat)
m.plot(LST_x, LST_y, 'bo', markersize=5,
       label='Lake Shore Traps')

SLB_x,SLB_y = m(NSVG_nsl['site_lon'].tolist()+SLB_Data['average_lon'].tolist(),
                  NSVG_nsl['site_lat'].tolist()+SLB_Data['average_lat'].tolist())
m.plot(SLB_x, SLB_y, 'ro', markersize=5,
       label='Schroeder-Lutsen Basalts')

MI_x,MI_y = m(SS_final_tc['average_lon'].tolist()+Palmer_data_trimmed['site_lon'].tolist(),
               SS_final_tc['average_lat'].tolist()+Palmer_data_trimmed['site_lat'].tolist())
m.plot(MI_x, MI_y, 'go', markersize=5,
       label='Michipicoten Island Formation')
plt.legend()
plt.show()
```



7 References

- Books, K., 1968, Magnetization of the lowermost Keweenawan lava flows in the Lake Superior area, Geological Survey research 1968, chapter D: U.S. Geological Survey Professional Paper, vol. P 0600-D, pp. 248–254.
- Books, K., 1972, Paleomagnetism of some Lake Superior Keweenawan rocks: U.S. Geological Survey Professional Paper, vol. P 0760, p. 42.
- Davis, D. and Green, J., 1997, Geochronology of the North American Midcontinent rift in western Lake Superior and implications for its geodynamic evolution: Canadian Journal of Earth Science, vol. 34, pp. 476–488, doi: 10.1139/e17-039.
- Diehl, J. and Haig, T., 1994, A paleomagnetic study of the lava flows within the Copper Harbour Conglomerate, Michigan: New results and implications: Canadian Journal of Earth Sciences, vol. 31, pp. 369–380, doi:10.1139/e94-034.
- Evans, D., 2009, The palaeomagnetically viable, long-lived and all-inclusive Rodinia supercontinent reconstruction: In Murphy, J., Keppie, J., and Hynes, A., eds., Ancient Orogens and Modern Analogues, Geological Society of London Special Publication, vol. 327, pp. 371–404, doi:10.1144/sp327.16.
- Fisher, N. I., Lewis, T., and Embleton, B. J. J., 1987, Statistical Analysis of Spherical Data: Cambridge University Press, doi:10.1017/CBO9780511623059.
- Kulakov, E. V., Smirnov, A. V., and Diehl, J. F., 2013, Paleomagnetism of ~1.09 Ga Lake Shore Traps (Keweenaw Peninsula, Michigan): new results and implications: Canadian Journal of Earth Sciences, vol. 50, pp. 1085–1096, doi:10.1139/cjes-2013-0003.
- Li, Z. X. et al., 2008, Assembly, configuration, and break-up history of Rodinia: A synthesis: Precambrian Research, vol. 160, pp. 179–210, doi:10.1016/j.precamres.2007.04.021.
- Malone, D. H., Stein, C. A., Craddock, J. P., Kley, J., Stein, S., and Malone, J. E., 2016, Maximum depositional age of the Neoproterozoic Jacobsville Sandstone, Michigan: Implications for the evolution of the Midcontinent Rift: Geosphere, doi:10.1130/GES01302.1, URL <http://geosphere.gsapubs.org/content/early/2016/07/12/GES01302.1.abstract>.
- McFadden, P. and McElhinny, M., 1990, Classification of the reversal test in palaeomagnetism: Geophysical Journal International, vol. 103, pp. 725–729.
- Palmer, H., 1970, Paleomagnetism and correlation of some Middle Keweenawan rocks, Lake Superior: Canadian Journal of Earth Science, vol. 7, pp. 1410–1436, doi:10.1139/e70-136.
- Palmer, H. and Davis, D., 1987, Paleomagnetism and U-Pb geochronology of volcanic rocks from Michipicoten Island, Lake Superior, Canada: precise calibration of the Keweenawan polar wander track: Precambrian Research, vol. 37, pp. 157–171, doi:10.1016/0301-9268(87)90077-5.
- Pisarevsky, S., Wingate, T., Powell, C. M., Johnson, S., and Evans, D., 2003, Models of Rodinia assembly and fragmentation: In Yoshida, M., Windley, B., and Dasgupta, S., eds., Proterozoic East Gondwana: Supercontinent Assembly and Breakup, The Geological Society of London Special Publications, vol. 206, pp. 35–55, doi:10.1144/gsl.sp.2003.206.01.04.
- Symons, D. T. A., Kawasaki, K., and Diehl, J. F., 2013, Age and genesis of the White Pine stratiform copper mineralization, northern Michigan, USA, from paleomagnetism: Geofluids, vol. 13, pp. 112–126, doi:10.1111/gfl.12024.
- Tauxe, L. and Kodama, K., 2009, Paleosecular variation models for ancient times: Clues from Keweenawan lava flows: Physics of the Earth and Planetary Interiors, vol. 177, pp. 31–45, doi:10.1016/j.pepi.2009.07.006.
- Tauxe, L., Shaar, R., Jonestrask, L., Swanson-Hysell, N., Minnett, R., Koppers, A., Constable, C., Jarboe, N., Gaastra, K., and Fairchild, L., 2016, PmagPy: Software package for paleomagnetic data analysis and a bridge to the Magnetics Information Consortium (MagIC) Database: Geochemistry, Geophysics, Geosystems, doi:10.1002/2016GC006307.
- Watson, G., 1956, A test for randomness of directions: Geophysical Journal International, vol. 7, pp. 160–161, doi:10.1111/j.1365-246X.1956.tb05561.x.