Pettinga et al., 2018, Morphometric scaling relationships in submarine channel–lobe systems: Geology, https://doi.org/10.1130/G45142.1.

Table DR1. Submarine Lobe Bodies and Associated Channel Dimensions

Fan system information								Submarine lobe bodies (LBs)									Submarine channels			Contextual information*			References
#	System name	Basin	Location	Fan age	Margin type	Confined/ unconfined	Fan type	Name	Hierarchy (H2-H4)	Length, L_{LB} (m)	Width, W_{LB} (m)	L_{LB}/W_{LB} (m/m)	Max thickness, H_{LB} (m)	Area, A_{LB} (km^2)	Volume, V_{LB} (km^3)	Width, W_{CH} (m)	Depth, H_{CH} (m)	Area, A_{CH} (m^2)	Drainage basin area (10^6 km^2)	Fluvial Q (km^3/yr)	Fluvial TSS (Mt/yr)	
1	Al Batha	Oman Basin, Arabian Sea	Offshore Oman	Last Glacial Maximum	passive	unconfined	basin floor	Al Batha Lobe Complex	4	95000	41000	2.3	30.0	1030.0	61.183 [C]	2000	130	204204 [C]	0.0055	–	–	Bourget et al., 2010
2	Al Batha	Oman Basin, Arabian Sea	Offshore Oman	Last Glacial Maximum	passive	unconfined	basin floor	Unit a	3	18715	22287	0.8	5.0	279.0	1.092 [C]	1000	20	15708 [C]	0.0055	–	–	Bourget et al., 2010
3	Al Batha	Oman Basin, Arabian Sea	Offshore Oman	Last Glacial Maximum	passive	unconfined	basin floor	Unit b	3	32321	23220	1.4	5.0	272.0	1.965 [C]	1000	20	15708 [C]	0.0055	–	–	Bourget et al., 2010
4	Al Batha	Oman Basin, Arabian Sea	Offshore Oman	Last Glacial Maximum	passive	unconfined	basin floor	Unit c	3	34702	12680	2.7	5.0	280.0	1.152 [C]	1000	20	15708 [C]	0.0055	–	–	Bourget et al., 2010
5	Al Batha	Oman Basin, Arabian Sea	Offshore Oman	Last Glacial Maximum	passive	unconfined	basin floor	Unit d	3	27595	17867	1.5	5.0	217.0	1.291 [C]	1000	20	15708 [C]	0.0055	–	–	Bourget et al., 2010
6	Al Batha	Oman Basin, Arabian Sea	Offshore Oman	Last Glacial Maximum	passive	unconfined	basin floor	"elementary body"	2	600	600	1.0	2.0	0.3 [C]	0.0004 [C]	–	–	–	0.0055	–	–	Bourget et al., 2010
7	Almeria	Alboran Sea	Offshore Spain	Late Pliocene/Quaternary	active	confined	basin floor	Almeria	4	45000	30000	1.5	–	1060.3 [C]	–	1000	53	41626 [C]	0.004	–	–	Garcia et al 2015; Alonso and Ercilla, 2003
8	Calahonda	Alboran Sea	Offshore Spain	Late Pliocene/Quaternary	active	unconfined	basin floor	Calahonda	4	12000	15000	0.8	–	141.4 [C]	–	–	–	–	0.0001	–	–	Garcia et al., 2015
9	Sacratif	Alboran Sea	Offshore Spain	Late Pliocene/Quaternary	active	confined	basin floor	Sacratif	4	24000	14000	1.7	–	263.9 [C]	–	–	–	–	0.0001	–	–	Garcia et al 2015; Alonso and Ercilla, 2003
10	Salobrena	Alboran Sea	Offshore Spain	Late Pliocene/Quaternary	active	confined	basin floor	Salobrena	4	17000	11000	1.5	–	146.9 [C]	–	–	–	–	0.001	–	–	Garcia et al., 2015
11	Fuengirola	Alboran Sea	Offshore Spain	Late Pliocene/Quaternary	active	unconfined	basin floor	Fuengirola	4	25000	20000	1.3	–	392.7 [C]	–	–	–	–	0.001	–	–	Garcia et al., 2015
12	Torrenueva	Alboran Sea	Offshore Spain	Late Pliocene/Quaternary	active	unconfined	basin floor	Torrenueva	4	12000	7000	1.7	–	66.0 [C]	–	–	–	–	0.0001	–	–	Garcia et al., 2015
13	Banos	Alboran Sea	Offshore Spain	Late Pliocene/Quaternary	active	unconfined	basin floor	Banos	4	16000	12000	1.3	–	150.8 [C]	–	–	–	–	0.0004	–	–	Garcia et al., 2015
14	Guadiaro	Alboran Sea	Offshore Spain	Late Pliocene/Quaternary	active	confined	basin floor	Guadiaro	4	14000	16000	0.9	–	175.9 [C]	–	300	38	8954 [C]	0.001	–	–	Garcia et al 2015; Alonso and Ercilla, 2003
15	La Linea	Alboran Sea	Offshore Spain	Late Pliocene/Quaternary	active	confined	basin floor	La Linea	4	5000	8000	0.6	–	31.4 [C]	–	–	–	–	0.0001	–	–	Garcia et al., 2015
16	Almirante Câmara	Campos Basin, Southern Atlantic	Offshore Brazil	Holocene	passive	confined	basin floor	Lobe (Lobo)	3	20000	10000	2.0	–	157.1 [C]	–	–	–	–	0.057	28	4	Machado et al., 2004
17	Amazon	Southern Atlantic	Offshore Brazil	Late Pleistocene	passive	unconfined	basin floor	A3	3	40000	22000	1.8	25.0	440.0	11.000	1224	18	17299 [C]	5.7	6400	1200	Juego et al., 2008
18	Amazon	Southern Atlantic	Offshore Brazil	Late Pleistocene	passive	unconfined	basin floor	A2	3	39000	11000	3.5	15.0	340.0	5.000	1224	18	17299 [C]	5.7	6400	1200	Juego et al., 2008
19	Amazon	Southern Atlantic	Offshore Brazil	Late Pleistocene	passive	unconfined	basin floor	A1	3	48000	17500	2.7	20.0	590.0	12.000	1224	18	17299 [C]	5.7	6400	1200	Juego et al., 2008
20	Amazon	Southern Atlantic	Offshore Brazil	Late Pleistocene	passive	unconfined	basin floor	1A	3	21000	6500	3.2	–	110.0	–	615	45	21736 [C]	5.7	6400	1200	Juego et al., 2008
21	Amazon	Southern Atlantic	Offshore Brazil	Late Pleistocene	passive	unconfined	basin floor	1C1	3	70000	9000	7.8	10.0	310.0	3.299 [C]	1080	55	46653 [C]	5.7	6400	1200	Juego et al., 2008
22	Amazon	Southern Atlantic	Offshore Brazil	Late Pleistocene	passive	unconfined	basin floor	1C2	3	80000	16000	5.0	10.0	655.0	6.702 [C]	1080	55	46653 [C]	5.7	6400	1200	Juego et al., 2008
23	Amazon	Southern Atlantic	Offshore Brazil	Late Pleistocene	passive	unconfined	basin floor	1C3	3	83000	20000	4.2	10.0	750.0	8.692 [C]	555	14	6103 [C]	5.7	6400	1200	Juego et al., 2008
24	Amazon	Southern Atlantic	Offshore Brazil	Late Pleistocene	passive	unconfined	basin floor	1C4	3	41000	11000	3.7	10.0	285.0	2.361 [C]	555	14	6103 [C]	5.7	6400	1200	Juego et al., 2008
25	Amazon	Southern Atlantic	Offshore Brazil	Late Pleistocene	passive	unconfined	basin floor	1Cx	3	60500	20000	3.0	10.0	680.0	6.336 [C]	555	14	6103 [C]	5.7	6400	1200	Juego et al., 2008
26	Amazon	Southern Atlantic	Offshore Brazil	Late Pleistocene	passive	unconfined	basin floor	1D-P	3	49000	18000	2.7	15.0	515.0	8.000	440	14	4838 [C]	5.7	6400	1200	Juego et al., 2008
27	Amazon	Southern Atlantic	Offshore Brazil	Late Pleistocene	passive	unconfined	basin floor	1D-D	3	65000	25000	2.6	15.0	1000.0	15.000	440	14	4838 [C]	5.7	6400	1200	Juego et al., 2008
28	Amazon	Southern Atlantic	Offshore Brazil	Late Pleistocene	passive	unconfined	basin floor	1E	3	49000	11000	4.5	10.0	485.0	4.800	790	18	11168 [C]	5.7	6400	1200	Juego et al., 2008
29	Amazon	Southern Atlantic	Offshore Brazil	Late Pleistocene	passive	unconfined	basin floor	1F	3	29500	19000	1.6	10.0	325.0	3.250	480	18	6786 [C]	5.7	6400	1200	Juego et al., 2008
30	Amazon	Southern Atlantic	Offshore Brazil	Late Pleistocene	passive	unconfined	basin floor	Br-P	3	45000	8500	5.3	10.0	220.0	2.200	740	24	13949 [C]	5.7	6400	1200	Juego et al., 2008
31	Amazon	Southern Atlantic	Offshore Brazil	Late Pleistocene	passive	unconfined	basin floor	Br-D	3	36000	12000	3.0	10.0	290.0	2.900	240	12	2262 [C]	5.7	6400	1200	Juego et al., 2008
32	Bering Sea Fans	Bering Sea	Offshore Alaska, USA	–	passive	unconfined	basin floor	Bering Turbidite System	4	235391	188365	1.2	–	29751.9	–	4000	10	31416 [C]	–	–	–	Kenyon and Millington, 1995
33	Bering Sea Fans	Bering Sea	Offshore Alaska, USA	–	passive	unconfined	basin floor	Umnak Turbidite System	4	427254	96435	4.4	–	24070.5	–	7000	100	549779 [C]	–	–	–	Kenyon and Millington, 1995
34	Bering Sea Fans	Bering Sea	Offshore Alaska, USA	–	passive	unconfined	basin floor	Pichinui Turbidite System	4	122862	53924	2.3	–	5026.5	–	5500	50	215984 [C]	–	–	–	Kenyon and Millington, 1995
35	Bering Sea Fans	Bering Sea	Offshore Alaska, USA	–	passive	unconfined	basin floor	B1	4	216026	38665	5.6	–	6512.9	–	–	–	–	–	–	–	Kenyon and Millington, 1995
36	Bering Sea Fans	Bering Sea	Offshore Alaska, USA	–	passive	unconfined	basin floor	B2	4	159172	36980	4.3	–	4623.0	–	–	–	–	–	–	–	Kenyon and Millington, 1995
37	Bering Sea Fans	Bering Sea	Offshore Alaska, USA	–	passive	unconfined	basin floor	B3	4	131151	29795	4.4	–	2869.0	–	–	–	–	–	–	–	Kenyon and Millington, 1995
38	Bering Sea Fans	Bering Sea	Offshore Alaska, USA	–	passive	unconfined	basin floor	B4	4	94952	35113	2.7	–	2502.4	–	–	–	–	–	–	–	Kenyon and Millington, 1995
39	Brazos-Trinity	Gulf of Mexico	Offshore Texas, USA	Late Pleistocene	passive	confined	ponded	Basin II lower sequence (DLC)	3	10000	8000	1.3	45.0	62.8 [C]	1.885 [C]	–	–	–	0.166	13	10.5	Beaubouef and Friedmann, 2000; Beaubouef et al., 2003
40	Brazos-Trinity	Gulf of Mexico	Offshore Texas, USA	Late Pleistocene	passive	confined	ponded	Basin II middle sequence (DLC)	3	11000	8000	1.4	30.0	69.1 [C]	1.382 [C]	–	–	–	0.166	13	10.5	Beaubouef and Friedmann, 2000; Beaubouef et al., 2003
41	Brazos-Trinity	Gulf of Mexico	Offshore Texas, USA	Late Pleistocene	passive	confined	ponded	Basin II upper sequence (DLC)	3	15000	12000	1.3	20.0	141.4 [C]	1.885 [C]	–	–	–	0.166	13	10.5	Beaubouef and Friedmann, 2000; Beaubouef et al., 2003
42	Brazos-Trinity	Gulf of Mexico	Offshore Texas, USA	Late Pleistocene	passive	confined	ponded	Basin IV lower fan (DLC)	3	10000	6000	1.7	37.0	47.1 [C]	1.162 [C]	–	–	–	0.166	13	10.5	Beaubouef and Friedmann, 2000; Beaubouef et al., 2003
43	Brazos-Trinity	Gulf of Mexico	Offshore Texas, USA	Late Pleistocene	passive	confined	ponded	Basin IV upper fan (DLC)	3	16000	9000	1.8	30.0	113.1 [C]	2.262 [C]	–	–	–	0.166	13	10.5	Beaubouef and Friedmann, 2000; Beaubouef et al., 2003
44	Brazos-Trinity	Gulf of Mexico	Offshore Texas, USA	Late Pleistocene	passive	confined	ponded	Basin II complex	4	15000	12000	1.3	150.0	134.0	14.137 [C]	–	–	–	0.166	13	10.5	Beaubouef and Friedmann, 2000; Beaubouef et al., 2003
45	Brazos-Trinity	Gulf of Mexico	Offshore Texas, USA	Late Pleistocene	passive	confined	ponded	Basin IV complex	4	16000	9000	1.8	200.0	112.0	15.080 [C]	–	–	–	0.166	13	10.5	Beaubouef and Friedmann, 2000; Beaubouef et al., 2003
46	Bute Inlet Fan	Bute inlet	British Columbia, Canada	Quaternary	passive	confined	basin floor	Lobe "Sand Splay"	3	9351	4418	2.1	–	37.9	–	246	24	4637 [C]	0.0076	11.7	5.7	Prior et al., 1986; Prior et al., 1987; Syvitski and Farrow, 1983
47	Celtic Fan	Bay of Biscay, Northern Atlantic	Offshore Ireland	Late Eocene	passive	unconfined	basin floor	Celtic Fan	4	215000	190000	1.1	–	30997.0	–	2200	50	86394 [C]	–	–	–	Zaragosi et al., 2003
48	Celtic Fan	Bay of Biscay, Northern Atlantic	Offshore Ireland	Late Eocene	passive	unconfined	basin floor	latest depositional lobe	3	115450	61171	1.9	–	4229.0	–	2200	50	86394 [C]	–	–	–	Zaragosi et al., 2003
49	Gendalo	Makassar Strait	Offshore Borneo, Indonesia	Upper Miocene	passive	unconfined	basin floor	Upper Gendalo1020 Lobe	3	7000	5000	1.4	–	27.5 [C]	–	–	–	–	–	–	–	Saller et al., 2008
50	Gendalo	Makassar Strait	Offshore Borneo, Indonesia	Upper Miocene	passive	unconfined	basin floor	Lower Gendalo1020 Lobe 1	3	4100	8100	0.5	–	26.1 [C]	–	–	–	–	–	–	–	Saller et al., 2008
51	Gendalo	Makassar Strait	Offshore Borneo, Indonesia	Upper Miocene	passive	unconfined	basin floor	Lower Gendalo1020 Lobe 2	3	2900	3900	0.7	–	8.9 [C]	–	–	–	–	–	–	–	Saller et al., 2008
52	Gendalo	Makassar Strait	Offshore Borneo, Indonesia	Upper Miocene	passive	unconfined	basin floor	Lower Gendalo1020 Lobe 3	3	3800	3000	1.3	–	9.0 [C]	–	–	–	–	–	–	–	Saller et al., 2008
53	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	A2	3	6001	4095	1.5	25.0	41.9	0.322 [C]	816	13	8332 [C]	0.001	0.5	0.002	Deptuck et al., 2008
54	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	A3	3	8704	6886	1.3	24.1	57.9	0.756 [C]	616	16	7741 [C]	0.001	0.5	0.002	Deptuck et al., 2008
55	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	B2	3	8210	5492	1.5	29.2	38.9	0.689 [C]	816	13	8332 [C]	0.001	0.5	0.002	Deptuck et al., 2008
56	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	B3	3	11695	5590	2.1	20.1	44.9	0.688 [C]	616	16	7741 [C]	0.001	0.5	0.002	Deptuck et al., 2008
57	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	B3a	2	11650	3200	3.6	20.0	29.3 [C]	0.390 [C]	816	13	8332 [C]	0.001	0.5	0.002	Deptuck et al., 2008
58	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	B3b	2	11680	3800	3.1	17.0	34.9 [C]	0.399 [C]	816	13	8332 [C]	0.001	0.5	0.002	Deptuck et al., 2008
59	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	C1	3	5312	4170	1.3	25.0	30.4	0.290 [C]	–	–	–	0.001	0.5	0.002	Deptuck et al., 2008
60	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	C1a	2	4950	3360	1.5	20.0	13.1 [C]	0.174 [C]	–	–	–	0.001	0.5	0.002	Deptuck et al., 2008
61	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	C1b	2	4950	3100	1.6	18.0	12.1 [C]	0.145 [C]	–	–	–	0.001	0.5	0.002	Deptuck et al., 2008
62	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	C2	3	12198	7305	1.7	27.3	53.0	1.274 [C]	816	13	8332 [C]	0.001	0.5	0.002	Deptuck et al., 2008
63	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	C3	3	13232	7683	1.7	40.8	69.9	2.172 [C]	616	16	7741 [C]	0.001	0.5	0.002	Deptuck et al., 2008
64	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	D3	3	14108	4915	2.9	37.5	47.0	1.362 [C]	616	16	7741 [C]	0.001	0.5	0.002	Deptuck et al., 2008
65	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	Fi	3	5889	3059	1.9	14.5	18.8	0.137 [C]	–	–	–	–	–	–	Deptuck et al., 2008
66	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	NG	3	5498	3925	1.4	19.5	17.0	0.220 [C]	816	13	8332 [C]	–	–	–	Deptuck et al., 2008
67	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	Pi	3	6365	2852	2.2	19.5	13.0	0.185 [C]	616	16	7741 [C]	–	–	–	Deptuck et al., 2008
68	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	x2	3	5461	2550	2.1	14.7	19.0	0.107 [C]	–	–	–	–	–	–	Deptuck et al., 2008
69	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	x3	3	1892	2884	0.7	12.8	6.3	0.037 [C]	–	–	–	–	–	–	Deptuck et al., 2008
70	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	x4	3	3001	2453	1.2	9.9	5.7	0.038 [C]	–	–	–	–	–	–	Deptuck et al., 2008
71	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	x5	3	2535	2151	1.2	9.1	4.0	0.026 [C]	–	–	–	–	–	–	Deptuck et al., 2008
72	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	x6	3	2917	1812	1.6	9.1	3.0	0.025 [C]	–	–	–	–	–	–	Deptuck et al., 2008
73	Golo Basin	Mediterranean Sea	Offshore East Corsica, France	Late Pleistocene	active	confined	basin floor	x7	3	2572	1284	2.0	8.9	4.2	0.015 [C]	–	–	–	–	–	–	Deptuck et al., 2008
74	Gulf of Cadiz, Aveiro	Gulf of Cadiz	Offshore Spain	Late Quaternary	active	confined	ponded	Aveiro Complex	4	6574	1575	4.2	10.0	5.1	0.054 [C]	250	2	393 [C]	–	–	–	Hanquiez et al., 2010
75	Gulf of Cadiz, Lolita	Gulf of Cadiz	Offshore Spain	Late Quaternary	active	confined	ponded	Lolita Complex	4	13059	2954	4.4	10.0	16.2	0.202 [C]	300	6	1414 [C]	–	–	–	Hanquiez et al., 2010
76	Gulf of Cadiz, Tasyo	Gulf of Cadiz	Offshore Spain	Late Quaternary	active	confined	ponded	Tasyo Complex	4	11419	3352	3.4	10.0	13.8	0.200 [C]	250	3	589 [C]	–	–	–	Hanquiez et al., 2010
77	Gulf of Cadiz, Aveiro	Gulf of Cadiz	Offshore Spain	Late Quaternary	active	confined	ponded	Aveiro Lobe A	3	2363	473	5.0	5.0	0.6	0.003 [C]	250	2	393 [C]	–	–	–	Hanquiez et al., 2010
78	Gulf of Cadiz, Aveiro	Gulf of Cadiz	Offshore Spain	Late Quaternary	active	confined	ponded	Aveiro Lobe B	3	5183	1016	5.1	5.0	2.4	0.014 [C]	250	2	393 [C]	–	–	–	Hanquiez et al., 2010
79	Gulf of Cadiz, Aveiro	Gulf of Cadiz	Offshore Spain	Late Quaternary	active	confined	ponded	Aveiro Lobe C	3	5440	1124	4.8	5.0	2.9	0.016 [C]	250	2	393 [C]	–	–	–	Hanquiez et al., 2010
80	Gulf of Cadiz, Lolita	Gulf of Cadiz	Offshore Spain	Late Quaternary	active	confined	ponded	Lolita Lobe A	3	5430	1263	4.3	5.0	3.9	0.018 [C]	300	6	1414 [C]	–	–	–	Hanquiez et al., 2010
81	Gulf of Cadiz, Lolita	Gulf of Cadiz	Offshore Spain	Late Quaternary	active	confined	ponded	Lolita Lobe B	3	6263	1641	3.8	5.0	7.2	0.027 [C]	300	6	1414 [C]	–	–	–	Hanquiez et al., 2010
82	Gulf of Cadiz, Lolita	Gulf of Cadiz	Offshore Spain	Late Quaternary	active	confined	ponded	Lolita Lobe C	3	10016	1994	5.0	5.0	6.5	0.052 [C]	300	6	1414 [C]	–	–	–	Hanquiez et al., 2010
83	Gulf of Cadiz, Tasyo	Gulf of Cadiz	Offshore Spain	Late Quaternary	active	confined	ponded	Tasyo Lobe A	3	11368	2279	5.0	5.0	8.6	0.068 [C]	250	3	589 [C]	–	–	–	Hanquiez et al., 2010
84	Gulf of Cadiz, Tasyo	Gulf of Cadiz	Offshore Spain	Late Quaternary	active	confined	ponded	Tasyo Lobe B	3	7462	1845	4.0	5.0	6.4	0.036 [C]	250	3	589 [C]	–	–	–	Hanquiez et al., 2010
85	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	Lower part of Basin floor fan	4	22000	22000	1.0	90.0	377.2	22.808 [C]	–	–	–	0.082	87	–	Saller et al., 2008
86	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	A	3	7000	5000	1.4	25.8	31.5	0.473 [C]	500	14.2	5576 [C]	0.082	87	–	Saller et al., 2008
87	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	B	3	5600	4000	1.4	33.9	19.1	0.398 [C]	400	16.8	5278 [C]	0.082	87	–	Saller et al., 2008
88	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	C	3	8000	5000	1.6	26.7	35.8	0.559 [C]	500	15.9	6244 [C]	0.082	87	–	Saller et al., 2008
89	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	D	3	2500	2500	1.0	42.0	9.8	0.137 [C]	250	8.4	1649 [C]	0.082	87	–	Saller et al., 2008
90	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	E	3	6400	6000	1.1	46.8	39.0	0.941 [C]	400	12.6	3958 [C]	0.082	87	–	Saller et al., 2008
91	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	F	3	7000	4700	1.5	42.8	38.9	0.737 [C]	300	10.9	2568 [C]	0.082	87	–	Saller et al., 2008
92	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	G	3	11000	5000	2.2	33.9	80.0	0.976 [C]	392	13.1	4033 [C]	0.082	87	–	Saller et al., 2008
93	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	H	3	7000	3000	2.3	34.7	33.6	0.382 [C]	392	13.1	4033 [C]	0.082	87	–	Saller et al., 2008
94	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	I	3	11000	7000	1.6	46.0	81.6	1.855 [C]	392	13.1	4033 [C]	0.082	87	–	Saller et al., 2008
95	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	J	3	13000	1800	7.2	33.1	28.2	0.406 [C]	–	–	–	0.082	87	–	Saller et al., 2008
96	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	K	3	4000	2500	1.6	31.5	29.7	0.165 [C]	392	13.1	4033 [C]	0.082	87	–	Saller et al., 2008
97	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	L	3	3700	1400	2.6	21.8	15.9	0.059 [C]	392	13.1	4033 [C]	0.082	87	–	Saller et al., 2008
98	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	M	3	12000	4500	2.7	34.7	51.0	0.981 [C]	392	13.1	4033 [C]	0.082	87	–	Saller et al., 2008
99	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	N	3	8400	4000	2.1	29.1	39.4	0.512 [C]	392	13.1	4033 [C]	0.082	87	–	Saller et al., 2008
100	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	O	3	4000	4000	1.0	29.1	8.2	0.244 [C]	392	13.1	4033 [C]	0.082	87	–	Saller et al., 2008
101	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	P	3	6000	2500	2.4	28.3	17.5	0.222 [C]	392	13.1	4033 [C]	0.082	87	–	Saller et al., 2008
102	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	Q	3	7000	2000	3.5	33.1	22.6	0.243 [C]	392	13.1	4033 [C]	0.082	87	–	Saller et al., 2008
103	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	R	3	5500	2700	2.0	35.5	20.7	0.276 [C]	392	13.1	4033 [C]	0.082	87	–	Saller et al., 2008
104	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	A1	2	800	650	1.2	10.1	0.4 [C]	0.003 [C]	161	12.6	1593 [C]	0.082	87	–	Saller et al., 2008
105	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	B1	2	1500	500	3.0	8.4	0.6 [C]	0.003 [C]	161	12.6	1593 [C]	0.082	87	–	Saller et al., 2008
106	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	C1	2	500	500	1.0	12.6	0.2 [C]	0.002 [C]	161	12.6	1593 [C]	0.082	87	–	Saller et al., 2008
107	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	C2	2	900	550	1.6	8.4	0.4 [C]	0.002 [C]	161	12.6	1593 [C]	0.082	87	–	Saller et al., 2008
108	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	C3	2	500	800	0.6	10.1	0.3 [C]	0.002 [C]	161	12.6	1593 [C]	0.082	87	–	Saller et al., 2008
109	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	C4	2	1000	850	1.2	6.7	0.7 [C]	0.003 [C]	161	12.6	1593 [C]	0.082	87	–	Saller et al., 2008
110	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	C5	2	900	400	2.3	5.9	0.3 [C]	0.001 [C]	161	12.6	1593 [C]	0.082	87	–	Saller et al., 2008
111	Kutai Basin	Makassar Strait	Offshore Borneo, Indonesia	Upper Pleistocene	active	confined	basin floor	C6	2	1200	800	1.5	10.1	0.8 [C]	0.005 [C]	161	12.6	1593 [C]	0.082	87	–	Saller et al., 2008
112

References Cited in Table DR1:

Alonso, B., and Ercilla, G., 2003, Small turbidite systems in a complex tectonic setting (SW Mediterranean Sea): Morphology and growth patterns: Marine and Petroleum Geology, v. 19, p. 1225–1240, doi: 10.1016/S0264-8172(03)00036-9.

Beaubouef, R.T., and Friedmann, S.J., 2000, High resolution seismic/sequence stratigraphic framework for the evolution of Pleistocene intra slope basins, western Gulf of Mexico: Depositional models and reservoir analogs, *in* Weimer, P., Slatt, R.M., Coleman, J., Rosen, N.C., Nelson, H., Bouma, A.H., Styzen, M.J., and Lawrence, D.T., eds., Deep-water reservoirs of the world: Gulf Coast Section SEPM 20th Annual Research Conference, p. 40–60, doi: 10.5724/gcs.00.15.0040.

Beaubouef, R.T., Abreu, V., and Adair, N.L., 2003, Ultra-high resolution 3-D characterization of deep-water deposits-I : A new approach to understanding the stratigraphic evolution of intra-slope depositional systems [abs.], AAPG Annual Convention Program, v. 12, p. A10.

Bourget, J., Zaragosi, S., Mulder, T., Schneider, J.L., Garlan, T., Van Toer, A., Mas, V., and Ellouz-Zimmermann, N., 2010, Hyperpycnal-fed turbidite lobe architecture and recent sedimentary processes: A case study from the Al Batha turbidite system, Oman margin: Sedimentary Geology, v. 229, p. 144–159, doi: 10.1016/j.sedgeo.2009.03.009.

Cossey, S., Kim, D.D., Yang, S., and Jung, H.Y., 2013, The identification and implication of injectites in the Shwe Gas Field, offshore northwestern Myanmar: AAPG Annual Convention and Exhibition, Pittsburgh, Pennsylvania, 19-22 May, 2013, AAPG Search and Discovery Article #20225,

Dennielou, B., et al., 2017, Morphology, structure, composition and build-up processes of the active channel-mouth lobe complex of the Congo deep-sea fan with inputs from remotely operated underwater vehicle (ROV) multibeam and video surveys: Deep-Sea Research Part II: Topical Studies in Oceanography, v. 142, p. 25–49, doi: 10.1016/j.dsr2.2017.03.010.

Deptuck, M.E., Piper, D.J.W., Savoye, B., and Gervais, A., 2008, Dimensions and architecture of late Pleistocene submarine lobes off the northern margin of East Corsica: Sedimentology, v. 55, p. 869–898, doi: 10.1111/j.1365-3091.2007.00926.x.

Droz, L., Kergoat, R., Cochonat, P., and Berné, S., 2001, Recent sedimentary events in the western Gulf of Lions (Western Mediterranean): Marine Geology, v. 176, p. 23–37, doi: 10.1016/S0025-3227(01)00147-5.

Droz, L., dos Reis, A.T., Rabineau, M., Berné, S., and Bellaiche, G., 2006, Quaternary turbidite systems on the northern margins of the Balearic Basin (Western Mediterranean): A synthesis: Geo-Marine Letters, v. 26, p. 347–359, doi: 10.1007/s00367-006-0044-0.

Fildani, A., and Normark, W.R., 2004, Late Quaternary evolution of channel and lobe complexes of Monterey Fan: Marine Geology, v. 206, p. 199–223, doi: 10.1016/j.margeo.2004.03.001.

García, M., Ercilla, G., Alonso, B., Estrada, F., Jané, G., Mena, A., Alvés, T., and Juan, C., 2015, Deep-water turbidite systems: a review of their elements, sedimentary processes and depositional models. Their characteristics on the Iberian margins: Boletín Geológico y Minero, v. 126, p. 189–218.

Hanquiez, V., Mulder, T., Toucanne, S., Lecroart, P., Bonnel, C., Marchès, E., and Gonthier, E., 2010, The sandy channel-lobe depositional systems in the Gulf of Cadiz: Gravity processes forced by contour current processes: Sedimentary Geology, v. 229, p. 110–123, doi: 10.1016/j.sedgeo.2009.05.008.

Jegou, I., Savoye, B., Pirmez, C., and Droz, L., 2008, Channel-mouth lobe complex of the recent Amazon Fan: The missing piece: Marine Geology, v. 252, p. 62–77, doi: 10.1016/j.margeo.2008.03.004.

Jegou, I., 2008, Etude de la transition chenal-levées/lobe dans les systèmes turbiditiques récents: Application à l'éventail turbiditique de l'Amazone et au Néofan du Petit-Rhône [Ph.D. thesis]: l'Université de Brest, 1-383 p.

Kenyon, N.H., and Millington, J., 1995, Contrasting deep-sea depositional systems in the Bering Sea, *in* Pickering, K.T., Hiscott, R.N., Kenyon, N.H., Ricci Lucchi, F., and Smith, R.D.A., eds., Atlas of Deep Water Environments: Architectural style in turbidite systems, London, Chapman & Hall, p. 196–202, doi: 10.1007/978-94-011-1234-5_28.

Kjøller, H.E., 2015, Source-to-sink system analysis of a modern submarine sedimentary system, Judd/Flett Sub-basins, west of Shetland. A possible analogue to ancient Paleocene subsurface sedimentary systems [M.S. thesis]: University of Copenhagen, 74 p.

Machado, L.C.R., Kowsmann, R.O., Almeida-Jr, W., Murakami, C.Y., Schreiner, S., Miller, D.J., and Piauilino, P.O.V., 2004, Geometria da porção proximal do sistema deposicional turbidítico moderno da Formação Carapebus, Bacia de Campos; modelo para heterogeneidades de reservatório Geometry: Boletim de Geociências da Petrobras, v. 12, p. 287–315.

Migeon, S., Ducassou, E., Le Gonidec, Y., Rouillard, P., Mascle, J., and Revel-Rolland, M., 2010, Lobe construction and sand/mud segregation by turbidity currents and debris flows on the western Nile deep-sea fan (Eastern Mediterranean): Sedimentary Geology, v. 229, p. 124–143, doi: 10.1016/j.sedgeo.2010.02.011.

Milliman, J.D., and Farnsworth, K.L., 2013, River discharge to the coastal ocean: a global synthesis: Cambridge, New York, Cambridge University Press, 394 p.

Normark, W.R., and Dickson, F.H., 1976a, Man-made turbidity currents in Lake Superior: Sedimentology, v. 23, p. 815–831, doi: 10.1111/j.1365-3091.1976.tb00110.x.

Normark, W.R., and Dickson, F.H., 1976b, Sublacustrine fan morphology in Lake Superior: AAPG Bulletin, v. 60, p. 1021–1036.

Normark, W.R., Piper, D.J.W., and Hess, G.R., 1979, Distributary channels, sand lobes, and mesotopography of Navy Submarine Fan, California Borderland, with applications to ancient fan sediments: Sedimentology, v. 26, p. 749–774, doi: 10.1002/9781444304473.ch36.

Picot, M., Droz, L., Marsset, T., Dennielou, B., and Bez, M., 2016, Controls on turbidite sedimentation: Insights from a quantitative approach of submarine channel and lobe architecture (Late Quaternary Congo Fan): Marine and Petroleum Geology, v. 72, p. 423–446, doi: 10.1016/j.marpetgeo.2016.02.004.

Prélat, A., Covault, J.A., Hodgson, D.M., Fildani, A., and Flint, S.S., 2010, Intrinsic controls on the range of volumes, morphologies, and dimensions of submarine lobes: Sedimentary Geology, v. 232, p. 66–76, doi: 10.1016/j.sedgeo.2010.09.010.

Prior, D.B., Bornhold, B.D., and Johns, M.W., 1986, Active sand transport along a fjord-bottom channel, Bute Inlet, British Columbia: Geology, v. 14, p. 581–584, doi: 10.1130/0091-7613(1986)14<581:ASTAAF>2.0.CO;2.

Prior, D.B., Bornhold, B.D., Wiseman, W.J., and Lowe, D.R., 1987, Turbidity current activity in a British Columbia fjord: Science, v. 237, p. 1330–1333, doi: 10.1126/science.237.4820.1330.

Saller, A., Werner, K., Sugiaman, F., Cebastiant, A., May, R., Glenn, D., and Barker, C., 2008, Characteristics of Pleistocene deep-water fan lobes and their application to an upper Miocene reservoir model, offshore East Kalimantan, Indonesia: AAPG Bulletin, v. 92, p. 919–949, doi: 10.1306/03310807110.

Stelting, C.E., Droz, L., Bouma, A.H., Coleman, J.M., Cremer, M., Meyer, A.W., Normark, W.R., Connell, S.O., and Stow, D.A. V, 1986, Late Pleistocene seismic stratigraphy of the Mississippi Fan, *in* Turner, K.L., ed., Initial Reports of the Deep Sea Drilling Project, v. 96, p. 40–55, doi: 10.2973/dsdp.proc.96.119.1986.

Syvitski, J.P.M., and Farrow, G.E., 1983, Structures and processes in bayhead deltas: Knight and Bute inlet, British Columbia: Sedimentary Geology, v. 36, p. 217–244, doi: 10.1016/0037-0738(83)90010-6.

Twichell, D.C., Kenyon, N.H., Parson, L.M., and McGregor, B.A., 1991, Depositional Patterns of the Mississippi Fan Surface: Evidence from GLORIA II and High-Resolution Seismic Profiles, *in* Weimer, P. and Link, M.H., eds., Seismic Facies and Sedimentary Processes of Submarine Fans and Turbidite Systems, New York, NY, Springer, p. 349–363, doi: 10.1007/978-1-4684-8276-8_19.

Twichell, D.C., Nelson, C.H., Kenyon, N.H., and Schwab, W.C., 2009, The influence of external Processes on the Holocene evolution of the Mississippi Fan, *in* Kneller, B., Martinsen, O.J., and McCaffrey, B., eds., External Controls on Deep-Water Depositional Systems, SEPM Special Publication No. 92, p. 145–157.

Wen, R., Sinding-Larsen, R., and Kenyon, N.H., 1995, Computer enhanced GLORIA sidescan sonar images of the surface of the Mississippi Fan, *in* Pickering, K.T., Hiscott, R.N., Kenyon, N.H., Ricci Lucchi, F., and Smith, R.D.A., eds., Atlas of Deep Water Environments: Architectural style in turbidite systems, London, Chapman & Hall, p. 297–299, doi: 10.1007/978-94-011-1234-5_44.

Wilde, P., Normark, W.R., and Chase, T.E., 1978, Channel sands and petroleum potential of Monterey deep-sea fan, California: AAPG Bulletin, v. 62, p. 967.

Yang, S.Y., and Kim, J.W., 2014, Pliocene basin-floor fan sedimentation in the Bay of Bengal (offshore northwest Myanmar): Marine and Petroleum Geology, v. 49, p. 45–58, doi: 10.1016/j.marpetgeo.2013.09.007.

Zaragosi, S., Le Suave, R., Bourillet, J.F., Auffret, G.A., and Garlan, T., 2003, Physiography of the Armorican turbidite system (Bay of Biscay), *in* Mienert, J. and Weaver, P., eds., European Margin Sediment Dynamics: Side-Scan Sonar and Seismic Images, Heidelberg, Germany, Springer, p. 245–248, doi: 10.1007/978-3-642-55846-7_40.

Zhang, J.J., Wu, S.H., Fan, T.E., Fan, H.J., Jiang, L., Chen, C., Wu, Q.Y., and Lin, P., 2016, Research on the architecture of submarine-fan lobes in the Niger Delta Basin, offshore West Africa: Journal of Palaeogeography, v. 5, p. 185–204, doi: 10.1016/j.jop.2016.05.005.

Figure DR1



References

[1] Picot, M., Droz, L., Marsset, T., Dennielou, B., and Bez, M., 2016, Controls on turbidite sedimentation: Insights from a quantitative approach of submarine channel and lobe architecture (Late Quaternary Congo Fan): Marine and Petroleum Geology, v. 72, p. 423–446, doi: 10.1016/j.marpetgeo.2016.02.004.

[2] Jegou, I., Savoye, B., Pirmez, C., and Droz, L., 2008, Channel-mouth lobe complex of the recent Amazon Fan: The missing piece: Marine Geology, v. 252, p. 62–77, doi: 10.1016/j.margeo.2008.03.004.

[3] Twichell, D.C., Kenyon, N.H., Parson, L.M., and McGregor, B.A., 1991, Depositional Patterns of the Mississippi Fan Surface: Evidence from GLORIA II and High-Resolution Seismic Profiles, in Weimer, P. and Link, M.H. eds., Seismic Facies and Sedimentary Processes of Submarine Fans and Turbidite Systems, New York, NY, Springer New York, p. 349–363, doi: 10.1007/978-1-4684-8276-8_19.

[4] Twichell, D.C., Nelson, C.H., Kenyon, N.H., and Schwab, W.C., 2009, The influence of external Processes on the Holocene evolution of the Mississippi Fan, in Kneller, B., Martinsen, O.J., and McCaffrey, B. eds., External Controls on Deep-Water Depositional Systems, SEPM Special Publication No. 92, p. 145–157.

[5] Droz, L., dos Reis, A.T., Rabineau, M., Berné, S., and Bellaiche, G., 2006, Quaternary turbidite systems on the northern margins of the Balearic Basin (Western Mediterranean): A synthesis: Geo-Marine Letters, v. 26, p. 347–359, doi: 10.1007/s00367-006-0044-0.

[5] Droz, L., dos Reis, A.T., Rabineau, M., Berné, S., and Bellaiche, G., 2006, Quaternary turbidite systems on the northern margins of the Balearic Basin (Western Mediterranean): A synthesis: Geo-Marine Letters, v. 26, p. 347–359, doi: 10.1007/s00367-006-0044-0.

[6] Jegou, I., 2008, Etude de la transition chenal-levées/lobe dans les systèmes turbiditiques récents: Application à l'éventail turbiditique de l'Amazone et au Néofan du Petit-Rhône [Ph.D. thesis]: l'Université de Brest, 1-383 p.

[7] Zaragosi, S., Auffret, G.A., Faugères, J.C., Garlan, T., Pujol, C., and Cortijo, E., 2000, Physiography and recent sediment distribution of the Celtic Deep-Sea Fan, Bay of Biscay: Marine Geology, v. 169, p. 207–237, doi: 10.1016/S0025-3227(00)00054-2.

{8] Bourget, J., Zaragosi, S., Mulder, T., Schneider, J.L., Garlan, T., Van Toer, A., Mas, V., and Ellouz-Zimmermann, N., 2010, Hyperpycnal-fed turbidite lobe architecture and recent sedimentary processes: A case study from the Al Batha turbidite system, Oman margin: Sedimentary Geology, v. 229, p. 144–159, doi: 10.1016/j.sedgeo.2009.03.009.

[9] Saller, A., Werner, K., Sugiaman, F., Cebastiant, A., May, R., Glenn, D., and Barker, C., 2008, Characteristics of Pleistocene deep-water fan lobes and their application to an upper Miocene reservoir model, offshore East Kalimantan, Indonesia: AAPG Bulletin, v. 92, p. 919–949, doi: 10.1306/03310807110.

[[10] Normark, W.R., Piper, D.J.W., and Hess, G.R., 1979, Distributary channels, sand lobes, and mesotopography of Navy Submarine Fan, California Borderland, with applications to ancient fan sediments: Sedimentology, v. 26, p. 749–774, doi: 10.1002/9781444304473.ch36.

[11] Deptuck, M.E., Piper, D.J.W., Savoye, B., and Gervais, A., 2008, Dimensions and architecture of late Pleistocene submarine lobes off the northern margin of East Corsica: Sedimentology, v. 55, p. 869–898, doi: 10.1111/j.1365-3091.2007.00926.x.

[12] Normark, W.R., and Dickson, F.H., 1976, Man-made turbidity currents in Lake Superior: Sedimentology, v. 23, p. 815–831, doi: 10.1111/j.1365-3091.1976.tb00110.x.

[13] Deptuck, M.E., Piper, D.J.W., Savoye, B., and Gervais, A., 2008, Dimensions and architecture of late Pleistocene submarine lobes off the northern margin of East Corsica: Sedimentology, v. 55, p. 869–898, doi:10.1111/j.1365-3091.2007.00926.x.